

# Beyond Cumulative Risk: A Dimensional Approach to Childhood Adversity

Katie A. McLaughlin<sup>1</sup> and Margaret A. Sheridan<sup>2</sup>

<sup>1</sup>Department of Psychology, University of Washington, and <sup>2</sup>Department of Psychology and Neuroscience, University of North Carolina at Chapel Hill

Current Directions in Psychological Science  
2016, Vol. 25(4) 239–245  
© The Author(s) 2016  
Reprints and permissions:  
sagepub.com/journalsPermissions.nav  
DOI: 10.1177/0963721416655883  
cdps.sagepub.com



## Abstract

Children who have experienced environmental adversity—such as abuse, neglect, or poverty—are more likely to develop physical and mental health problems, perform poorly at school, and have difficulties in social relationships than children who have not encountered adversity. What is less clear is how and why adverse early experiences exert such a profound influence on children’s development. Identifying developmental processes that are disrupted by adverse early environments is the key to developing better intervention strategies for children who have experienced adversity. Yet much existing research relies on a cumulative-risk approach that is unlikely to reveal these mechanisms. This approach tallies the number of distinct adversities experienced to create a risk score. This risk score fails to distinguish between distinct types of environmental experiences, implicitly assuming that very different experiences influence development through the same underlying mechanisms. We advance an alternative model. This novel approach conceptualizes adversity along distinct dimensions, emphasizes the central role of learning mechanisms, and distinguishes between different forms of adversity that might influence learning in distinct ways. A key advantage of this approach is that learning mechanisms provide clear targets for interventions aimed at preventing negative developmental outcomes in children who have experienced adversity.

## Keywords

childhood adversity, cumulative risk, stress, trauma, deprivation, learning, abuse, neglect, poverty

Few people dispute the notion that adverse environmental experiences in childhood—such as exposure to violence and chronic poverty—create a lasting imprint on emotion, cognition, behavior, and chances for success in adulthood. Children who have experienced adversity are more likely to develop psychopathology and chronic diseases, perform poorly at school, and have social and economic difficulties than children who have not encountered adversity (Felitti et al., 1998; Lansford et al., 2002; McLaughlin et al., 2012). What is less clear is how and why adverse early experiences exert such a profound influence on children’s development. Identifying the developmental processes that are disrupted by adverse early environments is the key to developing better intervention strategies to prevent the onset of problems in children who have experienced adversity.

Yet most current research is not designed in a way that can reveal mechanisms linking childhood adversity with developmental outcomes. We argue that the prevailing

approach for conceptualizing and measuring childhood adversity is, at best, not suited for studying mechanisms and may, at worst, obscure them. We first describe current approaches to studying childhood adversity and the dominant perspective on stress as the central mechanism linking these experiences with downstream outcomes. Next, we advance an alternative model for studying mechanisms linking childhood adversity with psychopathology and other developmental outcomes. This novel approach conceptualizes adversity along distinct dimensions, emphasizes the central role of learning mechanisms, and distinguishes between different forms of childhood adversity that might influence learning in distinct ways. A key

---

## Corresponding Author:

Katie McLaughlin, Department of Psychology, University of Washington, Box 351525, Seattle, WA 98195  
E-mail: mclaughk@uw.edu

advantage of our approach is that learning mechanisms provide clear targets for intervention.

### **The Prevailing Approach: Cumulative Risk**

Until recently, most research focused on single types of adversity, such as physical abuse, parental death, or poverty. A critical limitation of this approach is that it does not account for the fact that most children who have been exposed to one type of adversity have also experienced numerous others (Green et al., 2010; McLaughlin et al., 2012). When examining single types of adversity, it is impossible to determine whether a particular outcome (e.g., depression) is a consequence of the focal adversity of interest (e.g., physical abuse) or of other adversities the child experienced (e.g., poverty, neglect).

Recognition of the high co-occurrence of adversities led to a shift to the prevailing cumulative-risk approach. This approach tallies the number of adversities experienced to create a risk score (Evans, Li, & Whipple, 2013). For example, a child who experienced physical abuse, sexual abuse, and domestic violence would have a risk score of 3; a child who experienced poverty, neglect, and maternal depression would also have a risk score of 3. Cumulative risk thus focuses on the *number* of distinct adverse experiences rather than the severity or type of adversity (Evans et al., 2013).

The cumulative-risk approach has been widely adopted and has proved useful for highlighting the public-health importance of childhood adversity. Risk scores also can be used as a screening tool to identify children in greatest need of intervention. However, the cumulative-risk approach has significant limitations when used to identify mechanisms linking childhood adversity with developmental outcomes. Most notably, cumulative risk scores fail to distinguish between distinct types of environmental experience, implicitly assuming that all adverse experiences influence development through the same underlying mechanisms. In other words, risk scores assume that physical abuse, sexual abuse, and domestic violence influence children's development in exactly the same way as poverty, neglect, and maternal depression. This assumption is highly tenuous.


### ***Stress-response-system dysregulation as the central mechanism***

Is it possible that the vastly different social and environmental experiences encompassed by the construct of childhood adversity each influence development through the same underlying mechanisms? Advocates of cumulative risk argue that disruptions in the regulation of stress-response systems represent this common

mechanism (Evans et al., 2013). Specifically, allostatic load has been proposed as the process that explains how numerous forms of seemingly disparate adverse experiences influence the wide range of developmental outcomes associated with childhood adversity. The concept of allostatic load has been reviewed extensively elsewhere (McEwen, 2012; McEwen & Gianaros, 2010). Briefly, physiological regulatory systems—including the hypothalamic-pituitary-adrenal (HPA) axis and the autonomic nervous system (ANS)—respond to changing environmental demands, producing physiological changes that are adaptive in the short term but maladaptive in the long term. For example, chronic release of glucocorticoids leads to rapid improvements in immunity but maladaptive long-term changes in brain regions with high concentrations of glucocorticoid receptors, including the hippocampus, amygdala, and prefrontal cortex (McEwen, 2012). This long-term wear and tear resulting from chronic adaptation to stress is what is referred to as allostatic load. Extensive evidence suggests that adverse early environments disrupt stress-response system functioning (Gunnar & Quevedo, 2007). These disruptions are the central mechanism explaining downstream consequences of adversity in the cumulative-risk model.

Dysregulation in stress-response systems is clearly one pathway linking childhood adversity with developmental outcomes. But are stress pathways a universal mechanism? There are several problems with this assumption. First, associations of childhood adversity with stress-response-system functioning are inconsistent. Although numerous forms of adversity are associated with HPA-axis and ANS function, the specific nature of these associations varies widely across studies. The most commonly observed pattern involves blunted reactivity to environmental demands and globally reduced output (Gunnar & Vazquez, 2001; McLaughlin et al., 2015). However, numerous studies document the opposite pattern—elevated reactivity or globally increased output (Fries, Shirtcliff, & Pollak, 2008; Gunnar, Morison, Chisolm, & Schuder, 2001).

Second, stress-response-system dysregulation is inconsistently associated with developmental outcomes. Disruptions in stress-response systems are clearly involved in the onset of chronic physical health problems (Heim, Ehlert, & Helhammer, 2000). But they do not explain many disturbances in cognitive and social development commonly observed among children who have experienced adversity. For example, children exposed to neglect and poverty often have deficits in language abilities (Farah et al., 2006; Hildyard & Wolfe, 2002). There is no obvious link between stress-response-system dysregulation and language ability. Other mechanisms must be involved.



**Fig. 1.** A dimensional model of childhood adversity involving two central dimensions of threat and deprivation. Examples of commonly studied forms of adversity are placed along these dimensions based on the degree to which each experience typically involves threat and deprivation. Larger circles indicate greater variance in the degree to which the experience reflects the underlying dimension.

Finally, this pathway provides little in the way of intervention targets. How might we intervene to prevent the downstream consequences of childhood adversity based on the allostatic-load model, other than by attempting to prevent exposure to adversity in the first place? Although psychosocial interventions can influence cortisol regulation, the direction of these effects is remarkably inconsistent (Slopen, McLaughlin, & Shonkoff, 2014). Although the cumulative-risk approach has highlighted the importance of sensitive and responsive caregiving for adaptive development, it has provided few clues about how to intervene for children whose environments are characterized by an absence of such caregiving. Thus, although these models are useful in identifying children in need of intervention, they provide little guidance about *how* to intervene.

### A Novel Approach: Dimensions of Adversity

We have proposed an alternative model to facilitate the identification of developmental processes disrupted by childhood adversity other than the frequently invoked stress pathways (McLaughlin, Sheridan, & Lambert, 2014; Sheridan & McLaughlin, 2014). Identifying these mechanisms is critical for intervention development.

Our approach attempts to distill complex adverse experiences into core underlying dimensions that cut across multiple forms of adversity. Here, we focus on two specific dimensions of adversity whose influences on emotional, cognitive, and neurobiological development are at least partially distinct. Specifically, our model differentiates between experiences of *threat*—experiences involving harm or threat of harm, and *deprivation*—experiences involving an absence of expected inputs from the environment (see Fig. 1). Each dimension encompasses numerous experiences that reflect the underlying dimension to varying degrees. Experiences involving threat include observing community violence, witnessing domestic violence, and being the victim of chronic physical abuse. These experiences vary in the severity of threat involved, but all involve harm or threat of harm. Examples of deprivation include poverty, neglect, and institutional rearing, each of which involve an absence of expected cognitive inputs (e.g., complex language), social stimulation, and consistent interactions with adults to varying degrees.

Rather than counting the total number of adversities, our approach assesses the frequency and severity of experiences reflecting each dimension and examines them simultaneously in predicting developmental outcomes. This approach retains many benefits of cumulative

risk and provides additional advantages. Most notably, it can identify developmental mechanisms that are specific to certain dimensions of adversity and not others and determine whether such mechanisms vary in relation to severity of exposure.

### ***Beyond stress: The importance of learning***

Our model focuses on the central role of learning in explaining myriad developmental consequences of adversity not fully accounted for by stress pathways. Learning is a central process through which the environment shapes emotion, cognition, and behavior. Emerging research shows that learning processes are influenced by childhood adversity, that at least some of these associations vary across types of adversity, and that disruptions in learning are a mechanism in the link between adversity and developmental outcomes.

Emotional-learning processes, including fear and reward learning, are particularly important mechanisms in this regard. Associative-learning processes detect environmental cues associated with threat and reward and shape emotion, behavior, and neurobiological responses to those cues. Fear-learning processes rapidly detect potential threats and mobilize resources to respond (Delgado, Olsson, & Phelps, 2006). These processes govern acquisition of fear responses to stimuli associated with threat, extinction of fear following repeated stimulus encounters in the absence of threat, and competition between fear and extinction memories based on context. Reward-learning processes track the probability and magnitude of rewards associated with particular cues and influence feelings of pleasure during the anticipation and receipt of rewards as well as actions toward future rewards (Berridge & Kringelbach, 2008).

These forms of learning share basic neurobiological pathways with other associative learning processes such as pattern learning. Pattern learning refers to the detection of regularities in the environment when these regularities are not linked with rewards or punishments (e.g., that cue A always follows cue B). Pattern learning plays a central role in language acquisition and expertise (Romberg & Saffran, 2010; Spencer, Kaschak, Jones, & Lonigan, 2015).

Higher-order learning processes such as long-term memory and executive functions are also important candidate mechanisms explaining downstream consequences of adversity. In contrast to associative learning, which involves automatic processes present from early in development, long-term memory and executive functions are explicit learning skills for which effort and strategy can be employed to improve performance and that develop throughout childhood and adolescence.

### ***Childhood adversity influences learning***

Accumulating evidence shows that childhood adversity influences learning and that disruptions in learning processes are associated with downstream developmental outcomes, including psychopathology.

Exposure to threat is a core dimension of our model. We argue that disruptions in fear learning are an important mechanism through which threatening environments influence later development. Recent findings from our labs are consistent with this prediction, showing atypical fear conditioning among children who experienced environmental threats, including abuse and domestic violence. Children exposed to threat demonstrate poor discrimination of threat and safety cues during fear conditioning (McLaughlin et al., 2016). Whereas children without adversity exposure exhibit stronger fear responses to a stimulus paired with threat compared to one paired with safety, children exposed to threat exhibit fear responses of similar magnitude to threat and safety cues (McLaughlin et al., 2016), reflecting either generalization of fear to the safety cue or a generalized problem with associative learning. This pattern is specific to threat exposure, is not observed following deprivation, and explains the association of threat exposure with externalizing psychopathology.

Atypical reward learning has also been observed in children exposed to adversity, particularly those who experienced deprivation. In reward-learning tasks, typically developing children are faster and more accurate in responding to cues associated with high reward; children raised in deprived institutional settings fail to show this pattern (Sheridan, McLaughlin, Fox, Zeanah, & Nelson, under review). Disruptions in the neural circuitry that supports reward learning have been found in institutionally reared children (Mehta et al., 2010) and in adolescents who experienced neglect (Hanson, Hariri, & Williamson, 2015). It is possible that disruptions in reward learning emerge after other forms of adversity. Atypical reward learning and atypical neural responses to rewards have been found following child maltreatment (Dillon et al., 2009; Guyer et al., 2006). Because these studies did not distinguish between abuse and neglect, however, it is unknown whether reward learning is influenced only by deprivation or also by threat. Across studies, disruptions in reward processing and underlying neural circuitry have explained the link between deprivation and depression (Hanson et al., 2015; Sheridan et al., 2015).

Finally, disturbances in pattern learning, language, and executive functions have been found in children exposed to deprivation. Deficits in language and executive functions have been observed consistently among children raised in poverty (Noble, McCandliss, & Farah, 2007) and

in deprived institutional settings (Tibu et al., 2016; Windsor et al., 2011). These deficits explain the link between institutional rearing and attention-deficit/hyperactivity disorder (Tibu et al., 2016). Emerging evidence suggests that deprivation might also influence pattern learning (Sheridan et al., 2015) and long-term memory (Sheridan, How, Araujo, Schamberg, & Nelson, 2013), although greater work is needed in these areas.

### **Implications for intervention**

What does a learning perspective provide that stress models do not? Perhaps the most important advantage is that it provides clear targets for intervention. Behavioral interventions directly targeting emotional learning have been developed, evaluated, and refined for decades. These intervention techniques form the backbone of most empirically supported psychosocial treatments for anxiety, depression, behavior problems, and substance abuse in youth (Chorpita & Daleiden, 2009). For example, exposure-based interventions for anxiety facilitate extinction learning when that process does not occur naturally (Waters & Pine, 2016). Behavioral activation for depression aims to increase motivation for reward through repeated engagement in pleasurable activities. Behavioral interventions thus target the precise learning mechanisms that appear to be disrupted following childhood adversity. The development of interventions to improve memory and executive functions is a burgeoning area of research (Shipstead, Redick, & Engle, 2012).

Although behavioral interventions are effective in treating psychopathology, little research has examined their utility in preventing the onset of psychopathology in children who have experienced adversity. However, a recent study highlights the promise of behavioral interventions in this regard. A brief intervention providing behavioral-skill training to children who experienced traumatic violence prevented the onset of posttraumatic stress disorder and anxiety 3 months later (Berkowitz, Stover, & Marans, 2011). Greater research on the efficacy of brief behavioral interventions in preventing psychopathology following childhood adversity represents a critical next step for the field.

### **Conclusion**

There is little debate about the pervasive detrimental influence of childhood adversity on developmental outcomes. Similarly, broad consensus exists about the importance of developing effective interventions to prevent the downstream consequences of adversity. Although the prevailing cumulative-risk approach is useful for identifying children in need of intervention, it has done little to shed light on how to intervene. A dimensional approach focused on examining how specific types of adversity

influence learning and other mechanisms may hold greater promise in this regard. This approach is likely to reveal that some mechanisms are common across multiple dimensions of adversity and some are unique to particular experiences. Identifying these mechanisms and their specificity to particular forms of adversity will be critical for informing the development of efficient, effective interventions to prevent the negative developmental consequences of childhood adversity.

### **Recommended Reading**

- Evans, G. W., Li, D., & Whipple, S. S. (2013). Cumulative risk and child development. *Psychological Bulletin*, *139*, 1342–1396. A comprehensive review of the prevailing cumulative-risk approach—an approach distinct from the one we advance in this article.
- McEwen, B. S. (2012). (See References). An excellent and accessible review of the allostatic-load model.
- McLaughlin, K. A., Sheridan, M. A., Gold, A. L., Lambert, H. K., Heleniak, C., Duys, A., . . . Pine, D. S. (2016). (See References). Examines fear-learning mechanisms in children exposed to threatening early environments.
- McLaughlin, K. A., Sheridan, M. A., & Lambert, H. K. (2014). (See References). Provides a comprehensive review of the dimensional model of childhood adversity.

### **Declaration of Conflicting Interests**

The authors declared that they had no conflicts of interest with respect to their authorship or the publication of this article.

### **Funding**

This research was supported by National Institutes of Health Grants R01-MH103291 and R01-MH106482, an International Mental Health Research Organization Rising Star Award, and an Early Career Research Fellowship from the Jacobs Foundation to Katie A. McLaughlin.

### **References**

- Berkowitz, S. J., Stover, C. S., & Marans, S. R. (2011). The child and family traumatic stress intervention: Secondary prevention for youth at risk of developing PTSD. *Journal of Child Psychology and Psychiatry*, *52*, 676–685.
- Berridge, K. C., & Kringelbach, M. L. (2008). Affective neuroscience of pleasure: Reward in humans and animals. *Psychopharmacology*, *199*, 457–480.
- Chorpita, B. F., & Daleiden, E. L. (2009). Mapping evidence-based treatments for children and adolescents: Application of the distillation and matching model to 615 treatments from 322 randomized trials. *Journal of Consulting and Clinical Psychology*, *77*, 566–579.
- Delgado, M. R., Olsson, A., & Phelps, E. A. (2006). Extending animal models of fear conditioning to humans. *Biological Psychology*, *73*, 39–48.
- Dillon, D. G., Holmes, A. J., Birk, J. L., Brooks, N., Lyons-Ruth, K., & Pizzagalli, D. A. (2009). Childhood adversity is associated with left basal ganglia dysfunction during reward

- anticipation in adulthood. *Biological Psychiatry*, *66*, 206–213. doi:10.1016/j.biopsych.2009.02.019
- Evans, G. W., Li, D., & Whipple, S. S. (2013). Cumulative risk and child development. *Psychological Bulletin*, *139*, 1342–1396.
- Farah, M. J., Shera, D. M., Savage, J. H., Betancourt, L., Gianetta, J. M., Brodsky, N. L., . . . Hurt, H. (2006). Childhood poverty: Specific associations with neurocognitive development. *Brain Research*, *1110*, 166–174.
- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., . . . Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, *14*, 245–258.
- Fries, A. B. W., Shirtcliff, E. A., & Pollak, S. D. (2008). Neuroendocrine dysregulation following early social deprivation in children. *Developmental Psychobiology*, *50*, 588–599.
- Green, J. G., McLaughlin, K. A., Berglund, P., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2010). Childhood adversities and adult psychopathology in the National Comorbidity Survey Replication (NCS-R) I: Associations with first onset of DSM-IV disorders. *Archives of General Psychiatry*, *62*, 113–123.
- Gunnar, M. R., Morison, S. J., Chisolm, K., & Schuder, M. (2001). Salivary cortisol levels in children adopted from Romanian orphanages. *Development and Psychopathology*, *13*, 611–628.
- Gunnar, M. R., & Quevedo, K. (2007). The neurobiology of stress and development. *Annual Review of Psychology*, *58*, 145–173.
- Gunnar, M. R., & Vazquez, D. M. (2001). Low cortisol and a flattening of expected daytime rhythm: Potential indices of risk in human development. *Development and Psychopathology*, *13*, 515–538.
- Guyer, A. E., Kaufman, J., Hodgdon, H. B., Masten, C. L., Jazbec, S., Pine, D. S., & Ernst, M. (2006). Behavioral alterations in reward system function: The role of childhood maltreatment and psychopathology. *Journal of the American Academy of Child & Adolescent Psychiatry*, *45*, 1059–1067.
- Hanson, J. L., Hariri, A. R., & Williamson, D. E. (2015). Blunted ventral striatum development in adolescence reflects emotional neglect and predicts depressive symptoms. *Biological Psychiatry*, *78*, 598–605.
- Heim, C., Ehler, U., & Hellhammer, D. H. (2000). The potential role of hypocortisolism in the pathophysiology of stress-related bodily disorders. *Psychoneuroendocrinology*, *25*, 1–35.
- Hildyard, K. L., & Wolfe, D. A. (2002). Child neglect: Developmental issues and outcomes. *Child Abuse & Neglect*, *26*, 679–695.
- Lansford, J. E., Dodge, K. A., Petit, G. S., Bates, J. E., Crozier, J. C., & Kaplow, J. B. (2002). A 12-year prospective study of the long-term effects of early child physical maltreatment on psychological, behavioral, and academic problems in adolescence. *Archives of Pediatrics and Adolescent Medicine*, *156*, 824–830.
- McEwen, B. S. (2012). Brain on stress: How the social environment gets under the skin. *Proceedings of the National Academy of Sciences, USA*, *109*, 17180–17185.
- McEwen, B. S., & Gianaros, P. J. (2010). Central role of the brain in stress and adaptation: Links to socioeconomic status, health, and disease. *Annals of the New York Academy of Sciences*, *1186*, 190–222. doi:10.1111/j.1749-6632.2009.05331.x
- McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A., & Kessler, R. C. (2012). Childhood adversities and first onset of psychiatric disorders in a national sample of adolescents. *Archives of General Psychiatry*, *69*, 1151–1160.
- McLaughlin, K. A., Sheridan, M. A., Gold, A. L., Lambert, H. K., Helleniak, C., Duys, A., . . . Pine, D. S. (2016). Maltreatment exposure, brain structure, and fear conditioning in children. *Neuropsychopharmacology*, *41*, 1956–1964.
- McLaughlin, K. A., Sheridan, M. A., & Lambert, H. K. (2014). Childhood adversity and neural development: Deprivation and threat as distinct dimensions of early experience. *Neuroscience & Biobehavioral Reviews*, *47*, 578–591.
- McLaughlin, K. A., Sheridan, M. A., Tibu, F., Fox, N. A., Zeanah, C. H., & Nelson, C. A. (2015). Causal effects of the early caregiving environment on stress response system development in children. *Proceedings of the National Academy of Sciences, USA*, *112*, 5637–5642.
- Mehta, M. A., Gore-Langton, E., Golembo, N., Colvert, E., Williams, S. C. R., & Sonuga-Barke, E. (2010). Hyporesponsive reward anticipation in the basal ganglia following severe institutional deprivation early in life. *Journal of Cognitive Neuroscience*, *22*, 2316–2325. doi:10.1162/jocn.2009.21394
- Noble, K. G., McCandliss, B. D., & Farah, M. J. (2007). Socioeconomic gradients predict individual differences in neurocognitive abilities. *Developmental Science*, *10*, 464–480.
- Romberg, A. R., & Saffran, J. R. (2010). Statistical learning and language acquisition. *Wiley Interdisciplinary Reviews: Cognitive Science*, *1*, 906–914.
- Sheridan, M. A., How, J., Araujo, M., Schamberg, M. A., & Nelson, C. A. (2013). What are the links between maternal social status, hippocampal function, and HPA axis function in children? *Developmental Science*, *16*, 665–675.
- Sheridan, M. A., & McLaughlin, K. A. (2014). Dimensions of early experience and neural development: Deprivation and threat. *Trends in Cognitive Sciences*, *18*, 580–585.
- Sheridan, M. A., McLaughlin, K. A., Fox, N. A., Zeanah, C. H., & Nelson, C. A. (2015). *Early caregiving environment causally impacts reward response and neural structure, mediating risk for depression in adolescence*. Manuscript submitted for publication.
- Shipstead, Z., Redick, T. S., & Engle, R. W. (2012). Is working memory training effective? *Psychological Bulletin*, *138*, 628–654.
- Sloven, N., McLaughlin, K. A., & Shonkoff, J. P. (2014). Interventions to improve cortisol regulation in children: A systematic review. *Pediatrics*, *133*, 312–326.
- Spencer, M., Kaschak, M. P., Jones, J. L., & Lonigan, C. J. (2015). Statistical learning is related to early literacy-related skills. *Reading and Writing*, *28*, 467–490.
- Tibu, F., Sheridan, M. A., McLaughlin, K. A., Fox, N. A., Zeanah, C. H., & Nelson, C. A. (2016). Disruptions of working memory and inhibition mediate the association between

- exposure to institutionalization and symptoms of attention-deficit/hyperactivity disorder. *Psychological Medicine*, *46*, 529–541.
- Waters, A. M., & Pine, D. S. (2016). Evaluating differences in Pavlovian fear acquisition and extinction as predictors of outcome from cognitive behavioural therapy for anxious children. *Journal of Child Psychology and Psychiatry*. Advance online publication. doi: 10.1111/jcpp.12522
- Windsor, J., Benigno, J. P., Wing, C. A., Carroll, P. J., Koga, S. F., Nelson, C. A., . . . Zeanah, C. H. (2011). Effect of foster care on young children's language learning. *Child Development*, *82*, 1040–1046. doi:10.1111/j.1467-8624.2011.01604.x