

Published in final edited form as:

Trends Cogn Sci. 2014 November ; 18(11): 580–585. doi:10.1016/j.tics.2014.09.001.

Dimensions of Early Experience and Neural Development: Deprivation and Threat

Margaret A. Sheridan, Ph.D.^{#1} and Katie A. McLaughlin, Ph.D.^{#2}

¹Division of Developmental Medicine, Boston Children's Hospital, Harvard Medical School

²Department of Psychology, University of Washington

These authors contributed equally to this work.

Abstract

Over the past decade, a growing area of research has focused on adverse childhood experiences (ACEs) and their impacts on neural and developmental outcomes. Work in the field to-date has generally conceptualized ACEs in terms of exposure to stress while overlooking the underlying dimensions of environmental experience that may distinctly impact neural development. We propose a novel framework that differentiates between *deprivation* (absence of expected cognitive and social input) and *threat* (presence of a threat to one's physical integrity). We draw support for the neural basis of this distinction from studies on fear learning and sensory deprivation in animals to highlight potential mechanisms through which experiences of threat and deprivation could come to effect neural structure and function in humans.

The past decade has witnessed a proliferation of research on adverse childhood experiences and developmental outcomes. The term 'adverse childhood experience' (ACE) has been used to refer to a wide range of negative exposures during childhood that powerfully affect mental health, cognitive, and educational outcomes.¹ The strong relationship between ACEs and developmental outcomes has generated considerable interest in identifying the neurodevelopmental mechanisms that explain these associations. A small but rapidly growing body of work has examined the impact of ACEs on neural structure and function.^{2,3} However, prior work on this subject has made little attempt to identify the underlying dimensions of environmental experience that might influence neural development. We propose a novel conceptual framework for understanding the impact of ACEs on neural development that differentiates between experiences of *deprivation* and *threat*. Our intended contribution is to identify potential pathways through which deprivation and threat come to impact neural structure and function using basic neuroscience principles from animal

© 2014 Elsevier Ltd. All rights reserved.

The corresponding author is Margaret A. Sheridan. Contact information is as follows: Margaret A. Sheridan, Ph.D., Division of Developmental Medicine, Boston Children's Hospital, 1 Autumn Street Office 528, Boston, MA 02115, Phone: 857-218-5220, Margaret.Sheridan@childrens.harvard.edu; Katie McLaughlin, Ph.D., Department of Psychology, University of Washington, Box 351525, Seattle, WA 98195, Phone: 206-616-7863, mclaughk@uw.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

research. We highlight pathways beyond the most commonly hypothesized mechanism of stress exposure to suggest additional ways in which ACEs influence brain development. Our aim is not to comprehensively review existing evidence on ACEs and neural development in humans or animals, but to provide a conceptual framework to guide future research.

The long-term negative effects of ACEs on developmental outcomes have been documented for decades. This research historically focused on single types of adversity, such as abuse and neglect. Recent studies have examined associations between *number* of ACEs and developmental outcomes,⁴ based on evidence that different types of ACEs frequently co-occur.¹ An unintended consequence of this approach has been an oversimplification of the boundaries between distinct types of environmental experience. One example of this problem involves use of the term “early-life stress” (ELS), which is used to refer to disparate experiences ranging from institutionalization to maternal depression and marital conflict,^{5,6} and obscures differences between these experiences that are likely to have important implications for understanding their impact on neural development. Characterizing underlying dimensions of environmental experience associated with diverse forms of adversity is critical for identifying their distinct effects on neural development, an essential first step in identifying mechanisms linking ACEs to developmental outcomes.

Here we propose a novel conceptual framework for studying the effects of ACEs on neural development. The central distinction we make is between experiences of *deprivation* and *threat* (Table 1). We suggest that these dimensions differentially influence neurodevelopment. We do not propose that exposure to deprivation and threat occur independently for children, as many ACEs co-occur. Instead we propose that they can be measured separately (Figure 1) and have unique effects on neurodevelopment. Below we separately describe deprivation and threat. Within each section, we first review mechanisms of neural development from animal neuroscience and describe how deprivation and threat influence these mechanisms. Next, we highlight emerging work in humans examining the neural consequences of ACEs. We end by proposing directions for future research that will help to determine the utility of our proposed framework.

The contribution of this perspective within the larger literature on ACEs and neurodevelopment is to highlight the importance of conceptualizing and measuring underlying dimensions of environmental experience reflected in frequently studied exposures, such as abuse, neglect, and poverty, because those dimensions may differentially influence neural outcomes. Critically, because fine-grained measurement of these dimensions has not been undertaken in human studies of neurodevelopment and because prior studies have focused on specific types of exposure (e.g., abuse) often without measuring or reporting co-occurring exposures (e.g., neglect), any conclusions regarding the consistency of existing human work with our proposed framework are necessarily tentative. Moreover, some exposures inherently involve high degrees of both deprivation and threat. For example, institutionalization involves the complete absence of an attachment figure in early development,⁷ an experience that involves not only deprivation in expected inputs but also can represent a significant threat to survival for an infant. Importantly, we do not suggest that deprivation and threat are the only dimensions of experience that are important or that all ACEs can be conceptualized solely along these dimensions. Rather, we propose

that these are two dimensions of experience that have not previously been differentiated with regard to their distinct influences on neural development.

Deprivation

The dimension of deprivation refers to the absence of species- or age-expectant environmental inputs, specifically a lack of expected cognitive and social inputs. We argue that the animal neuroscience literature examining the effects of sensory deprivation on sensory cortex development can be used as a model for understanding the neural consequences of deprivation in complex cognitive and social inputs in humans. Specifically, we suggest that an early environment without cognitive enrichment will yield a neural structure designed to deal with low complexity environments. Thus exposure to cognitive and social deprivation in children would result in: a) age-specific reductions in thickness and volume of association cortex, due in part to early over-pruning of synaptic connections, lower numbers of synaptic connections, and reduced dendritic branching; and b) reduced performance on tasks that depend on these areas (e.g., complex cognitive tasks). Reductions in cortical thickness should be most pronounced in regions of association cortex that are recruited for processing complex cognitive and social inputs, including prefrontal cortex (PFC), superior and inferior parietal cortex, and superior temporal cortex.

Insights from animal models

A central principal of neuroscience, developed and elaborated over the last half century, is that early experience shapes brain structure and function through pruning of synaptic connections in the central nervous system (CNS). Decades of work have documented that CNS development contains two distinct phases of synaptic growth: proliferation and pruning. Synaptic proliferation occurs in a period beginning during the third trimester, peaking three months after birth, and ending before the second year of life.^{8,9} During this period, there are rapid increases in the ratio of asymmetrical to symmetrical synapses, synaptic density, and total number of synapses.^{10,11} Following proliferation, a period of pruning of synaptic connections occurs and continues for an extended period through childhood and adolescence. In humans, synaptic elimination occurs earlier for primary sensory cortex and later for association cortex, although the final density of synapses in adulthood across cortical areas is not different.^{8,10} Pruning is dependent on co-activation: as two cells co-activate, the association between them strengthens, trophic factors are transmitted, and it becomes more likely that the connection will persist. The emergent system reflects the relative effectiveness of various pathways, theoretically yielding the most efficient system for the environment in it developed.

Early work on the effect of experience on neural development examined the effect of visual deprivation on visual cortex structure and function through direct manipulation of visual input. This work documents that visual deprivation early in development leads to irreversible changes in the structure and function of primary visual cortex resulting from a radical reduction in synapses (see Figure 2 for details). A second literature has investigated the impact of more general deprivation exposure. Global deprivation due to single rodent housing early in development—associated with decreased visual, auditory, and social inputs—results in widespread decreases in dendritic arborization, spines, neuronal depth and

cortical thickness.¹²⁻¹⁵ These changes are at least partially reversible through exposure to enriching, cognitively-stimulating environments following deprivation.¹⁴ In sum, there is evidence from the animal literature that decreases in environmental input during development within a single modality (e.g., vision) decreases dendritic arborization and number of synapses in corresponding sensory cortex and across multiple modalities (e.g., global deprivation) results in similar changes that are widespread throughout the cortex.

Insights from human studies

Given observed changes in cortical structure following sensory deprivation in rodents, cognitive and social deprivation in humans is likely to result in reductions in cortical thickness. These reductions are likely to be most pronounced in association cortex, because these regions develop and are influenced by environmental inputs for the longest period of time, and result in deficits in cognitive and social functions reliant on these regions of association cortex (e.g., language, executive function, spatial navigation, social cognition). As noted above, it is difficult to garner conclusive evidence for this hypothesis from the current literature given the way that ACEs have been measured and reported in existing studies. However, studies of children reared in institutional settings or exposed to neglect provide an opportunity to examine patterns of neural structure and function following exposure to environments that are clearly characterized by deprivation. In these environments most children will experience reduced and low quality cognitive (e.g., language exposure) and social (e.g., caregiver interactions) inputs.^{16,17} Studies examining neural structure in children exposed to these types of deprived environments observe reductions in cortical thickness in association cortex,^{18,19} disruptions in PFC function,²⁰ and declines in associated cognitive functions.^{21,22} Other environments, such as poverty, are not inherently characterized by deprivation, (i.e., it is possible to be poor and to have typical exposure to cognitive and social inputs) but serve as a *marker* for a greater likelihood of deprivation in exposure to enriching and cognitively complex environments.²³ Poverty has also been associated with reductions in cortical thickness in PFC,^{24,25} disruptions in PFC function,^{26,27} and declines in associated cognitive functions.^{28,29}

Threat

In contrast to the proposed impact of deprivation, we suggest that early threat exposure is associated with changes in neural circuits that underlie emotional learning, including the hippocampus, amygdala, and ventromedial PFC (vmPFC). Specifically, early threat exposure is associated with: a) changes in adult hippocampal morphology and function, including reduced dendritic spines and arborization and poor hippocampal function in learning and memory tasks; b) changes in amygdala function due to novel pairing of threat cues with previously neutral stimuli and heightened salience of emotional information, resulting in elevated amygdala activation to emotional stimuli and increased vigilance and attention to threat-related cues; and c) under-recruitment of vmPFC due to stronger representation of conditioned fear than extinction memories, resulting in reduced vmPFC thickness, low vmPFC recruitment during extinction recall and other types of emotional processing, and low structural and functional connectivity of vmPFC with amygdala and hippocampus.

Insights from Animal Models

Fear is a defensive mechanism that promotes survival. The neural circuitry underlying fear learning is well-characterized in animals; detailed reviews can be found elsewhere.^{30,31} Fear learning occurs when a previously innocuous stimulus is paired with an aversive stimulus, such that the neutral stimulus comes to elicit the behavioral and neurobiological responses associated with the aversive stimulus. This learning happens automatically, allowing threats to quickly elicit defensive responses. The amygdala is necessary for acquisition and expression of conditioned fear, and the hippocampus is involved in fear acquisition for complex stimuli.³² Learned fear generally abates with the passage of time as a result of extinction learning, although extinguished fear can be re-activated through a variety of processes, including exposure to novel threats.³³ Successful fear extinction involves recruitment of the vmPFC, which is required for retrieval of extinction learning, resulting in inhibition of the amygdala and dampened fear expression.³⁴

Exposure to threats early in development alters neural circuitry underlying fear learning. Here we focus on fear-eliciting paradigms, including shock, restraint, predator odor, and minimal bedding, which results in erratic, rough maternal care.³⁵ We do not review maternal separation paradigms, because they conflate dimensions of threat and deprivation. Early threat exposure results in lasting changes in hippocampus and amygdala structure and function. Specifically, early threat leads to reduced dendritic length and branching in adulthood, blunted long-term potentiation in the hippocampus, and impairments in hippocampus-dependent learning and memory,³⁶⁻⁴⁰ including impaired contextual fear learning and extinction of context-dependent fear.⁴¹⁻⁴³ Early threat also predicts dendritic atrophy in vmPFC and poor synaptic transmission between vmPFC and hippocampus.⁴³ In the amygdala, early threat is associated with increased dendritic spines, elevated basal activity and response to novel or stressful tasks, deficits in inhibitory pathways regulating amygdala activity, and widespread changes in gene expression.⁴³⁻⁴⁶

Insights from human studies

In rodents, early exposure to threat results in stable changes in the neural systems underlying fear learning. Thus, in humans, we expect that early threat exposure will be associated with parallel changes in fear learning and in the structure and function of the neural systems that support emotional learning: the hippocampus, amygdala, and vmPFC. Existing studies of children exposed to threatening environments, including physical and sexual abuse, domestic violence, and other violent trauma—which share the characteristic of being significant threats to survival—provide an opportunity to evaluate associations of threat with neural circuits underlying fear learning. Consistent with animal findings, early threat exposure is associated with: a) reduced hippocampal volume in adulthood but not childhood;^{47,48} b) poor hippocampal function on learning and memory tasks;⁴⁹ c) elevated amygdala reactivity to threatening stimuli (e.g., angry faces);^{50,51} d) attention biases that facilitate the identification of threats;⁵² e) reduced vmPFC volume,^{53,54} and f) reduced resting-state amygdala-vmPFC connectivity.⁵⁵ Importantly, threat exposure often co-occurs with deprivation (e.g., neglect), making it impossible to conclude that these patterns are specifically the result of threat.

Recommendations for Future Research

The exposures that give rise to experiences of deprivation and threat co-occur at high rates in children and adolescents.¹ This co-occurrence has generated many of the methodological and conceptual challenges in identifying dimensions of experience that influence specific aspects of neural development. One way of testing the pathways proposed here is to identify children who experience only one form of adversity. However, finding such a sample is not only difficult, but also would not accurately represent the population of children exposed to ACEs. Instead, we propose that future studies examining neural development in children exposed to ACEs should measure these underlying dimensions of experience, in addition to the traditional categories of exposure, to determine whether deprivation and threat are indeed uniquely associated with the patterns of neural development proposed here. We have focused on two dimensions of experience that are particularly likely to impact neural development, but there are undoubtedly others (e.g., chaos or unpredictability and loss). Future studies should identify other key dimensions of experience and characterize their impact on the developing brain.

Despite consistencies in the neural circuitry underlying fear learning in animals and humans, there is also a surprising lack of human research on how early threat influences fear learning across development. This represents a critical area for future research. Future studies should also attempt experimental manipulation of specific aspects of experience (e.g., increasing enriching cognitive experiences for institutionalized children) to identify the causal pathways through which environmental experience shapes neurodevelopment.

Concluding Remarks

We propose a novel conceptual framework for understanding the impact of ACEs on neural development. Our approach argues that the field must move beyond the prevailing approach of examining the impact of complex and co-occurring exposures on brain development to distilling those complex experiences into their core underlying dimensions. Two important dimensions that appear to have distinct effects on neural development are deprivation and threat. Existing evidence from human studies is consistent with animal work examining how deprivation and threat influence neural development, although additional work is needed to determine the utility of our proposed framework. We believe that such an approach will improve our understanding of how atypical experience influences the developing brain and, ultimately, confers risk for adverse developmental outcomes.

Acknowledgments

This research was supported by grants from the National Institutes of Health (K01-MH092555 to Sheridan and K01-MH092526 to McLaughlin).

References

1. McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky A, Kessler RC. Childhood adversities and first onset of psychiatric disorders in a national sample of adolescents. *Archives of General Psychiatry*. 2012; 69:1151–1160. [PubMed: 23117636]

2. Hart H, Rubia K. Neuroimaging of child abuse: a critical review. *Frontiers in Human Neuroscience*. 2012; 6 Article 52.
3. Hackman DA, Farah MJ. Socioeconomic status and the developing brain. *Trends in Cognitive Sciences*. 2009; 13:65–73. [PubMed: 19135405]
4. Schilling EA, Aseltine RH, Gore S. The impact of cumulative childhood adversity on young adult mental health: Measures, models, and interpretations. *Social Science and Medicine*. 2008; 66:1140–1151. [PubMed: 18177989]
5. Burghy CA, Stodola DE, Ruttelle PL, et al. Developmental pathways to amygdala-prefrontal function and internalizing symptoms in adolescence. *Nature Neuroscience*. 2012; 15:1736–1741.
6. Cohen MM, Jing D, Yang RR, Tottenham N, Lee FS, Casey BJ. Early-life stress has persistent effects on amygdala function and development in mice and humans. *Proceedings of the National Academy of Sciences*. 2013; 110:18274–18278.
7. Tottenham N. Human amygdala development in the absence of species-typical expected caregiving. *Developmental Psychobiology*. 2012; 54:598–611. [PubMed: 22714586]
8. Huttenlocher PR, Dabholkar AS. Regional differences in synaptogenesis in human cerebral cortex. *Journal of Comprehensive Neurology*. 1997; 387:167–178.
9. Petanjek Z, Judas M, Simic G, et al. Extraordinary neoteny of synaptic spines in the human prefrontal cortex. *Proceedings of the National Academy of Sciences*. 2011; 108(32):13281–13286. 2011/07/25/.
10. Huttenlocher PR, de Courten C. The development of synapses in striate cortex of man. *Hum Neurobiol*. 1987; 6(1):1–9. 1987. [PubMed: 3583840]
11. Rakic P, Bourgeois JP, Eckenhoff MF, Zecevic N, Goldman-Rakic PS. Concurrent overproduction of synapses in diverse regions of the primate cerebral cortex. *Science (New York, N.Y.)*. 1986; 232(4747):232–235. 1986/04/11/.
12. Bennett EL, Rosenzweig MR, Diamond MC, Morimoto H, Hebert M. Effects of successive environments on brain measures. *Physiol. Behav*. 1974; 12(4):621–631. 1974/04//. [PubMed: 4824387]
13. Diamond MC, Law F, Rhodes H, et al. Increases in cortical depth and glia numbers in rats subjected to enriched environment. *J. Comp. Neurol*. 1966; 128(1):117–126. 1966/09//. [PubMed: 4165855]
14. Diamond MC, Rosenzweig MR, Bennett EL, Lindner B, Lyon L. Effects of environmental enrichment and impoverishment on rat cerebral cortex. *J. Neurobiol*. 1972; 3(1):47–64. 1972. [PubMed: 5028293]
15. Globus A, Rosenzweig MR, Bennett EL, Diamond MC. Effects of differential experience on dendritic spine counts in rat cerebral cortex. *J Comp Physiol Psychol*. 1973; 82(2):175–181. 1973/02//. [PubMed: 4571892]
16. Smyke AT, Koga S, Johnson DE, et al. The caregiving context in institution-reared and family-reared infants and toddlers in Romania. *Journal of Child Psychology and Psychiatry*. 2007; 48:210–218. [PubMed: 17300560]
17. Hart, B.; Risley, TR. Meaningful differences in the everyday experiences of young American children. Paul H Brooks Publishing; Baltimore: 1995.
18. McLaughlin KA, Sheridan MA, Winter W, Fox NA, Zeanah CH, Nelson CA. Widespread reductions in cortical thickness following severe early-life deprivation: A neurodevelopmental pathway to ADHD. *Biological Psychiatry*. in press.
19. Sheridan MA, Fox NA, Zeanah CH, McLaughlin KA, Nelson CA. Variation in neural development as a result of exposure to institutionalization early in childhood. *Proceedings of the National Academy of Sciences*. 2012; 109(32):12927–12932.
20. Chugani HT, Behen ME, Muzik O, Juhász C, Nagy F, Chugani DC. Local brain functional activity following early deprivation: a study of postinstitutionalized Romanian orphans. *NeuroImage*. 2001; 14(6):1290–1301. 2001/12//. [PubMed: 11707085]
21. Beckett C, Castle J, Rutter M, Sonuga-Barke EJS. VI. Institutional deprivation, specific cognitive functions, and scholastic achievement: English and Romanian Adoptee (ERA) study findings. *Monographs of the Society for Research in Child Development*. 2010; 75(1):125–142. 2010/04//. [PubMed: 20500636]

22. Bos KJ, Fox NA, Zeanah CH, Nelson CA. Effects of early psychosocial deprivation on the development of memory and executive function. *Frontiers in Behavioral Neuroscience*. 2009; 3:16. [PubMed: 19750200]
23. Bradley RH, Conyn RF, Burchinal M, McAdoo HP, Coll CG. The home environments of children in the United States part II: relations with behavioral development through age thirteen. *Child development*. 2001; 72(6):1868–1886. 2001/12/Nov- undefined. [PubMed: 11768150]
24. Hanson JL, Hair N, Shen DG, et al. Family poverty affects the rate of human infant brain growth. *PLoS one*. 2013; 8(12) 2013.
25. Noble KG, Houston SM, Kan E, Sowell ER. Neural correlates of socioeconomic status in the developing human brain. *Developmental science*. 2012; 15(4):516–527. 2012/07//. [PubMed: 22709401]
26. Raizada RDS, Richards TL, Meltzoff A, Kuhl PK. Socioeconomic status predicts hemispheric specialization of the left inferior frontal gyrus in young children. *NeuroImage*. 2008; 40(3):1392–1401. 2008/04/15/. [PubMed: 18308588]
27. Sheridan MA, Sarsour K, Jutte D, D’Esposito M, Boyce WT. The impact of social disparity on prefrontal function in childhood. *PLoS One*. 2012; 7(4):e35744. [PubMed: 22563395]
28. Bradley RH, Corwyn RF, McAdoo HP, Coll CG. The home environments of children in the United States part I: variations by age, ethnicity, and poverty status. *Child development*. 2001; 72(6): 1844–1867. 2001/12/Nov- undefined. [PubMed: 11768149]
29. Linver MR, Brooks-Gunn J, Kohen DE. Family processes as pathways from income to young children’s development. *Developmental psychology*. 2002; 38(5):719–734. 2002/09//. [PubMed: 12220050]
30. Kim JJ, Jung MW. Neural circuits and mechanisms involved in Pavlovian fear conditioning: A critical review. *Neuroscience and Biobehavioral Reviews*. 2006; 30:188–202. [PubMed: 16120461]
31. Johansen JP, Cain CK, Ostroff LE, LeDoux JE. Molecular mechanisms of fear learning and memory. *Cell*. 2011; 147:509–524. [PubMed: 22036561]
32. Phillips RG, LeDoux JE. Differential contributions of amygdala and hippocampus to cued and contextual fear conditioning. *Behavioral Neuroscience*. 1992; 106:274–285. [PubMed: 1590953]
33. Bouton ME. Context, ambiguity, and unlearning: Sources of relapse after behavioral extinction. *Biological Psychiatry*. 2002; 52:976–986. [PubMed: 12437938]
34. Quirk GJ, Russo GK, Barron JL, Lebron K. The role of ventromedial prefrontal cortex in the recovery of extinguished fear. *Journal of Neuroscience*. 2000; 20:6225–6231. [PubMed: 10934272]
35. Roth TL, Sullivan RM. Memory of early maltreatment: Neonatal behavioral and neural correlates of maternal maltreatment within the context of classical conditioning. *Biological Psychiatry*. 2005; 57:823–831. [PubMed: 15820702]
36. Eiland L, Ramroop J, Hill MN, Manley J, McEwen BS. Chronic juvenile stress produces corticolimbic dendritic architectural remodeling and modulates emotional behavior in male and female rats. *Psychoneuroendocrinology*. 2012; 37:39–47. [PubMed: 21658845]
37. Rice CJ, Sandman CA, Lenjavi MR, Baram TZ. A novel mouse model for acute and long-lasting consequences of early life stress. *Endocrinology*. 2008; 149:4892–4900. [PubMed: 18566122]
38. Brunson KL, Kramar E, Lin B, et al. Mechanisms of late-onset cognitive decline after early-life stress. *Journal of Neuroscience*. 2005; 25:9328–9338. [PubMed: 16221841]
39. Ivy AS, Rex CS, Chen Y, et al. Hippocampal dysfunction and cognitive impairments provoked by chronic early-life stress involve excessive activation of CRH receptors. *Journal of Neuroscience*. 2010; 30:13005–13015. [PubMed: 20881118]
40. Isgor C, Kabbaj M, Akil H, Watson SJ. Delayed effects of chronic variable stress during peripubertal-juvenile period of hippocampal morphology and on cognitive and stress axis function in rats. *Hippocampus*. 2004; 14:636–648. [PubMed: 15301440]
41. Matsumoto M, Togashi H, Konno K, et al. Early postnatal stress alters the extinction of context-dependent conditioned fear in adult rats. *Pharmacology, Biochemistry and Behavior*. 2008; 89:247–252.

42. Toledo-Rodriguez M, Sandi C. Stress before puberty exerts a sex- and age-related impact on auditory and contextual fear conditioning in the rat. *Neural Plasticity*. 2007; 2007 Article ID 71203.
43. Eiland L, McEwen BS. Early life stress followed by subsequent adult chronic stress potentiates anxiety and blunts hippocampal structural remodeling. *Hippocampus*. 2012; 22:82–91. [PubMed: 20848608]
44. Sarro EC, Sullivan RM, Barr G. Unpredictable neonatal stress enhances adult anxiety and alters amygdala gene expression related to serotonin and GABA. *Neuroscience*. in press.
45. Sevelinges Y, Mouly A-M, Raineke C, Moriceau S, Forest C, Sullivan RM. Adult depression-like behavior, amygdala and olfactory cortex functions are restored by odor previously paired with shock during infancy. *Developmental Cognitive Neuroscience*. 2011; 1:77–87. [PubMed: 21037982]
46. Raineke C, Rincon Cortes M, Belnoue L, Sullivan RM. Effects of early-life abuse differ across development: Infant social behavior deficits are followed adolescent depressive-like behaviors mediated by the amygdala. *Journal of Neuroscience*. 2012; 32:7758–7765. [PubMed: 22649253]
47. Teicher MH, Anderson CM, Polcari A. Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proceedings of the National Academy of Sciences*. 2012; 109:E563–572.
48. Andersen SL, Teicher MH. Stress, sensitive periods and maturational events in adolescent depression. *Trends in Neuroscience*. 2008; 31:183–191.
49. Carrion VG, Haas BW, Garrett A, Song S, Reiss AL. Reduced hippocampal activity in youth with posttraumatic stress symptoms: An fMRI study. *Journal of Pediatric Psychology*. 2010; 35:559–569. [PubMed: 19995868]
50. McCrory EJ, De Brito SA, Sebastian CL, et al. Heightened neural reactivity to threat in child victims of family violence. *Current Biology*. 2011; 21:R947–948. [PubMed: 22153160]
51. McCrory EJ, De Brito SA, Kelly PA, et al. Amygdala activation in maltreated children during pre-attentive emotional processing. *British Journal of Psychiatry*. 2013; 202:269–276. [PubMed: 23470285]
52. Pollak SD, Sinha P. Effects of early experience on children’s recognition of facial displays of emotion. *Development and Psychopathology*. 2002; 38:784–791.
53. Hanson JL, Chung MK, Avants BB, et al. Early stress is associated with alterations in the orbitofrontal cortex: A tensor-based morphometry investigation of brain structure and behavioral risk. *Journal of Neuroscience*. 2010; 30:7466–7472. [PubMed: 20519521]
54. De Brito SA, Viding E, Sebastian CL, et al. Reduced orbitofrontal and temporal gray matter in a community sample of maltreated children. *Journal of Child Psychology and Psychiatry*. 2013; 54:105–112. [PubMed: 22880630]
55. Herringa RJ, Birn RM, Ruttle PL, Stodola DE, Davidson RJ, Essex MJ. Childhood maltreatment is associated with altered fear circuitry and increased internalizing symptoms by late adolescence. *Proceedings of the National Academy of Sciences*. 2013; 110:19119–19124.

Highlights

- New framework differentiates between depriving and threatening adverse childhood experiences
- Deprivation, or the absence of expected inputs, impacts proliferation and pruning
- Threat, or the presence of atypical trauma experiences, impacts fear learning
- These patterns can be observed in humans and animals





Figure 1. Dimensions of threat and deprivation associated with commonly occurring adverse childhood experiences (ACEs)

¹ Importantly, we argue that threat and deprivation are *dimensions* of experience that can be measured among children exposed to a wide ranges of ACEs, both those that occur in isolation (e.g., a single incident of community violence exposure) and those that are co-occurring (e.g., physical abuse and physical neglect). We use the term complex exposures to refer to experiences that in most cases involve aspects of both threat and deprivation.

Mechanism of Deprivation



A. Number of synapses in layer I, for two regions in area 17, corresponding to central and peripheral representations of the binocular visual field
N70: Normal 70 days; N220: Normal 220 days; DR220: Dark reared 220 days of maturity.

B. Ratio of the mean volume in the early blind subjects relative sighted controls at each voxel. The color bar displays the percentage difference in average volume between groups from greater than 30% to less than 30%.

Early work identifying the effect of deprivation on neural structure and function comes from investigations of the effect of sensory deprivation. A. This figure shows decreased synapses visual cortex of kittens raised in complete visual deprivation, or “dark rearing” (DR220) from post-natal day 70 – 220. This decrease in synapses is accompanied by a decrease in thickness compared to typically reared kittens (N220; reproduced with permission from O’Kusky, 1985a). **B.** Similarly, in humans, congenital blindness is associated with less volume in visual cortex compared to sighted or late-blind participants (reproduced with permission from Leporé et al., 2010).

Figure 2. The effects of early visual deprivation on cortical structure in animals and humans

Table 1
Definitions of Threat and Deprivation

Threat	Deprivation
Experiences of threat involve the presence of an atypical (i.e., unexpected) experience characterized by actual or threatened death, injury, sexual violation, or other harm to one’s physical integrity. Our definition of threat is consistent with the definition of a traumatic event in the Diagnostic and Statistical Manual of Mental Disorders – Fifth Edition (DSM-5). ¹	Experiences of deprivation involve the absence of expected environmental inputs in cognitive (e.g., language) and social domains as well as the absence of species- and age-typical complexity in environmental stimulation.

¹ American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.