

Low Emotional Awareness as a Transdiagnostic Mechanism Underlying Psychopathology in
Adolescence

David G. Weissman¹

Erik C. Nook¹

Aridenne A. Dews¹

Adam Bryant Miller²

Hilary K. Lambert³

Stephanie F. Sasse¹

Leah H. Somerville¹

Katie A. McLaughlin¹

¹Department of Psychology, Harvard University, Cambridge, MA

²Department of Psychology and Neuroscience, University of North Carolina, Chapel Hill, NC

³Department of Psychology, University of Washington, Seattle, WA

Abstract

The ability to identify and label one's emotions is a precursor to effective emotion regulation, suggesting that emotional awareness is important for mental health. We evaluated how emotional awareness was related to psychopathology and whether low emotional awareness was a transdiagnostic mechanism explaining the increase in psychopathology during the transition to adolescence and as a function of childhood trauma—specifically violence exposure. In Study 1, children and adolescents (N=120, aged 7-19 years) reported on emotional awareness and psychopathology. Emotional awareness was negatively associated with psychopathology (p-factor) and decreased across age in females but not males. In Study 2 (N=262, aged 8-16 years), we replicated these findings and demonstrated longitudinally that low emotional awareness mediated increases in p-factor as a function of age in females and violence exposure. These findings indicate that low emotional awareness may be a transdiagnostic mechanism linking adolescent development, sex, and trauma with the emergence of psychopathology.

Emotional awareness reflects one's subjective understanding of their emotional state (Lane & Schwartz, 1987; Lumley, Neely, & Burger, 2007). Emotional awareness is central to the experience of discrete emotions (Barrett, 2006) and is thought to be a prerequisite to many forms of effective emotion regulation (Barrett, Gross, Christensen, & Benvenuto, 2001; Kalokerinos, Erbas, Ceulemans, & Kuppens, 2019). Low emotional awareness is associated with many forms of psychopathology (Deborde, Vanwalleghem Maury, & Aitel, 2015; Frewen, Dozois, Neufeld, & Lanius, 2008; Hendryx, Haviland, & Shaw, 1991), suggesting that it may represent a transdiagnostic factor associated with increased vulnerability for psychopathology. We investigate this possibility in the current study. We additionally evaluate whether low emotional awareness is a mechanism that contributes to other well-established patterns of risk for transdiagnostic psychopathology. Specifically, we determine whether low emotional awareness explains increases in transdiagnostic psychopathology during the transition to adolescence and as a function of exposure to trauma in childhood—specifically, interpersonal violence exposure. Greater insight into these transdiagnostic mechanisms may inform the development of more effective early interventions.

The construct of emotional awareness can be understood through the lens of the constructionist model of emotion, which posits that individuals use emotion concepts to “construct” specific emotional experiences from an instance of core affect (i.e., one's momentary neurophysiological state that emerges within an environmental context; Barrett, 2006, 2017). Thus, emotional awareness requires perceiving one's internal bodily state—a process called interoception—and conceptualizing it in terms of an emotion label that best unites the current context with past experience (Satpute & Lindquist, 2019). This construction of an emotional experience then guides regulatory responses to the individual's current context (Kashdan,

Barrett, & McKnight, 2015). As such, factors that give rise to individual differences in interoception or emotion conceptualization across development may influence emotional awareness, emotion regulation, and ultimately psychopathology (Murphy, Brewer, Catmur, & Bird, 2017).

Although one might posit that emotional awareness should improve across adolescence as individuals master increasingly sophisticated emotional and non-emotional language (Nook, Sasse, Lambert, McLaughlin, & Somerville, 2017), adolescents actually exhibit greater difficulty differentiating their emotions compared to both younger and older ages (Nook, Sasse, Lambert, McLaughlin, & Somerville, 2018). These opposing forces impacting emotion conceptualization across age (i.e., increasing verbal abilities vs. decreasing ease at identifying emotional states) could generate either increased or decreased emotional awareness. In fact, studies on the development of emotional awareness have produced inconsistent results: Some studies show increased emotional awareness across ages 11-18 (Gatta et al., 2014; Säkkinen, Kaltiala-Heino, Ranta, Haataja, & Joukamaa, 2007), but others find decreased emotional awareness with pubertal development (van der Crujisen, Murphy, & Bird, 2019). If emotional awareness decreases during the transition to adolescence, it might be one factor explaining the well-established increase in risk for psychopathology that emerges transdiagnostically during this period (Kessler et al., 2005).

In addition to these developmental patterns, exposure to childhood adversity—particularly trauma exposure—is a potent risk factor for multiple forms of psychopathology (Green et al., 2010; McLaughlin et al., 2012), as well as for the transdiagnostic general psychopathology factor (i.e., p-factor) (Caspi et al., 2014; Weissman et al., 2019). Trauma exposure may contribute to lower emotional awareness through impacts on both interoception

and emotion conceptualization. Exposure to extreme or uncontrollable stressors during childhood—like exposure to violence and other forms of trauma—has been associated with blunted autonomic nervous system and cortisol reactivity to stressors as well as disruptions in diurnal cortisol patterns, typically with low morning levels and a shallow decline across the day (Carpenter, Shattuck, Tyrka, Geraciotti, & Price, 2011; Gordis, Feres, Olezeski, Rabkin, & Trickett, 2010; Gunnar & Vazquez, 2001; Harkness, Stewart, & Wynne-Edwards, 2011; Heleniak, McLaughlin, Ormel, & Riese, 2016; MacMillan et al., 2009; McLaughlin, Sheridan, Alves, & Mendes, 2014; Trickett, Noll, Susman, Shenk, & Putnam, 2010). These changes to physiological stress response systems may weaken the coupling between subjective experiences of negative emotion and corresponding physiological reactivity, ultimately manifesting as low emotional awareness.

Indeed, childhood adversity, including violence exposure, is associated with problems identifying and differentiating between emotions (Pears & Fisher, 2005; Shenk, Putnam, & Noll, 2013), which may be attributable to less frequent use of emotional words and poorer scaffolding of emotional concepts and experiences by parents (Denham, Mitchell-Copeland, Strandberg, Auerbach, & Blair, 1997; Pollak, Cicchetti, Hornung, & Reed, 2000; Salmon, O’Kearney, Reese, & Fortune, 2016). Children exposed to violence become particularly attuned to the detection of anger at the expense of differentiating between other negative emotions (Pollak et al., 2000; Pollak & Sinha, 2002; Shenk et al., 2013). Thus, violence exposure may contribute to reduced sensitivity to changes in one’s own bodily state as well as a more limited emotional vocabulary and less differentiated emotion concepts. Consistent with this possibility, exposure to stressful life events in adolescence prospectively predicts declines in emotional awareness over time (McLaughlin & Hatzenbuehler, 2009; McLaughlin, Hatzenbuehler, & Hilt, 2009).

Existing evidence suggests that low emotional awareness may be a factor that contributes to the association of multiple risk factors with transdiagnostic psychopathology. Indeed, low emotional awareness is associated with many forms of psychopathology, including substance abuse (Rybakowski, Ziółkowski, Zasadzka, & Brzeziński, 1988; Taylor, Parker, & Bagby, 1990), conduct disorder (Deborde et al., 2015), depression (Honkalampi, Hintikka, Laukkanen, & Viinamäki, 2002), anxiety (Hendryx et al., 1991), PTSD (Frewen et al., 2008), obsessive compulsive disorder (Grabe et al., 2006), and schizophrenia (van 't Wout, Aleman, Bermond, & Kahn, 2007). Two potential pathways could explain these associations. First, low emotional awareness may disrupt the ability to regulate emotions (Barrett et al., 2001; Kalokerinos et al., 2019). Second, because low emotional awareness is associated with more difficulty recognizing emotions in others (Grynberg et al., 2012), low emotional awareness could impair social functioning and in turn contribute to multiple forms of psychopathology (Collin, Bindra, Raju, Gillberg, & Minnis, 2013). Both of these pathways could explain why low emotional awareness may be a transdiagnostic mechanism conveying risk for multiple forms of psychopathology emerging in middle childhood and adolescence.

In line with this perspective, recent conceptualizations of psychopathology have shifted away from categorical diagnoses and instead advocated for focusing on underlying dimensions that reflect broader psychopathology. Empirical evidence supports the existence of a transdiagnostic psychopathology factor, termed the general factor or p-factor, that reflects the co-occurrence of psychopathology symptoms in children, adolescents, and adults (Caspi et al., 2014; Castellanos-Ryan et al., 2016; Gomez, Stavropoulos, Vance, & Griffiths, 2018; Laceulle, Vollebergh, & Ormel, 2015; Lahey et al., 2012; Martel et al., 2017; Murray, Eisner, & Ribeaud, 2016; Neumann et al., 2016; Patalay et al., 2015; Schaefer et al., 2018; Snyder, Young, &

Hankin, 2017). Adolescence is a period of risk for the onset of most forms of psychopathology (Kessler et al., 2005), and considerable evidence supports the notion that childhood violence exposure is associated with transdiagnostic risk for psychopathology (Caspi et al., 2014; Keyes et al., 2012; Weissman et al., 2019). However, the mechanisms that underlie this transdiagnostic risk have only recently been examined. In prior work, we demonstrated that elevations in emotional reactivity, engagement in maladaptive emotion regulation strategies (e.g., rumination), and altered patterns of neural response in the brain's salience network to threat cues are mechanisms linking childhood violence exposure with the emergence of psychopathology over time (Weissman et al., 2019; Weissman et al., in press). Here, we investigate whether low emotional awareness may be an additional mechanism linking both adolescent development and violence exposure with the p-factor.

We investigated factors that influence emotional awareness in childhood and adolescence and the association between self-reported emotional awareness and the p-factor in two studies. In Study 1 we modeled the structure of psychopathology in a community sample of children and adolescents and evaluated the age-related differences in p-factor and self-reported emotional awareness, and the association between emotional awareness the p-factor concurrently. In Study 2, we extended this work by examining the influence of both age and childhood violence exposure on emotional awareness and the p-factor both concurrently and prospectively over time. Finally, we investigated whether low emotional awareness was a mechanism underlying the association of age and violence exposure with increases in the p-factor over time. Given prior work suggesting that the developmental trajectory of emotional awareness varies in males and females (van der Crujisen et al., 2019), we also investigated whether sex was related to emotional awareness or moderated associations of age and violence exposure with emotional

awareness in both studies.

Study 1

Method

Participants. Data were drawn from a cross-sectional study of emotional development (Nook, Sasse, Lambert, McLaughlin, & Somerville, 2017, 2018; Nook et al., in press) conducted in a community sample spanning ages 4 to 25 years. Psychopathology measures were only validated for participants aged 7-19 years. Thus, our analytic sample was restricted to the 120 participants (62 female) in this age range. All participants were native English speakers who were compensated for their time and recruited from communities surrounding Harvard University and the University of Washington. Participants provided informed written consent/assent, and minor participants received written permission for their participation from a parent or legal guardian. The Committee on the Use of Human Subjects at Harvard University and the University of Washington Institutional Review Board approved all research procedures.

Emotional awareness. Low emotional awareness was measured using 12 items (e.g. “I have feelings I can’t quite identify”) from the Alexithymia Questionnaire for Children, excluding the 8 items of the externally oriented thinking subscale, which has low reliability in children and adolescents (Rieffe, Oosterveld, & Terwogt, 2006). Items are rated on a 5-point Likert scale from “Strongly Disagree” to “Strongly Agree.” This scale reliably measures low emotional awareness in children and adolescents (Heaven, Ciarrochi, & Hurrell, 2010; Loas, Braun, Delhayé, & Linkowski, 2017) and had good internal consistency in this sample ($\alpha = .89$). Higher scores indicate lower emotional awareness. Emotional awareness measures were available for 104 of the 120 participants.

Symptoms of psychopathology. Depression symptoms were assessed with the Children's Depression Inventory-2 (CDI-2), a recently revised version of the widely used self-report measure of depressive symptoms in children and adolescents (Kovacs, 1992, 2011). The CDI has demonstrated good reliability and validity among children and adolescents (Craighead, Smucker, Craighead, & Ilardi, 1998), and it demonstrated excellent internal consistency in our sample ($\alpha = .90$). Anxiety symptoms were assessed with the Screen for Child Anxiety Related Emotional Disorders (SCARED), which measures anxiety disorder symptoms across five domains: panic/somatic, generalized anxiety, separation anxiety, social phobia, and school phobia (Birmaher et al., 1997). The SCARED has sound psychometric properties (Birmaher et al., 1999, 1997) and excellent internal consistency in our sample ($\alpha = .93$). Externalizing symptoms were assessed using both child and caregiver reports on the Youth Self-Report (YSR) and Child Behavior Checklist (CBCL) (Achenbach, 1991). The YSR/CBCL scales are among the most widely used measures of youth emotional and behavioral problems. The higher raw score between the CBCL and YSR was used from the Attention Problems, Rule Breaking Behaviors, and Aggressive Behavior subscales. Symptoms of PTSD were assessed using child- and parent-report versions of the UCLA PTSD Reaction Index (PTSD-RI) (Steinberg, Brymer, Decker, & Pynoos, 2004). The PTSD-RI assesses PTSD re-experiencing, avoidance/numbing, and hyperarousal symptoms according to DSM-IV criteria. A total symptom severity score is generated by summing all items. The PTSD-RI has sound psychometric properties (Steinberg et al., 2013), and had excellent internal consistency in our sample ($\alpha = .92$). The higher of the parent and child reported PTSD symptom severity score was used.

Defining p-factor. Following Caspi et al. (2014) and recent replications (Laceulle et al., 2015; Schaefer et al., 2018; Weissman et al., 2019), we performed confirmatory factor analysis

(CFA) to test two standard models: a correlated-factors model specifying Internalizing and Externalizing latent factors and a bi-factor model specifying both a General Psychopathology latent factor (p-factor) and residual Internalizing and Externalizing factors. In order to ensure that our latent factors were not being driven by one or more indicators simply because of measurement differences across psychopathology instruments (i.e., different number of items, scoring, etc.), we binned scores on each indicator into deciles prior to CFA analyses. All CFA analyses were performed in MPlus version 8.1 (Muthén & Muthén, 2012). Given that our observed indicator variables were slightly skewed and kurtotic, we used the robust maximum likelihood estimator (MLR), which employs a sandwich estimator to arrive at standard errors robust to non-normality of observations. MLR performs well in modest sample sizes with skewed data, as in the present study (Li, 2016). We assessed the relative fit of each model using the Akaike Information Criterion (AIC), Bayesian Information Criterion (BIC) and the Sample Adjusted BIC.

Analysis. Linear regression was used to evaluate associations of age, sex, and their interaction with low emotional awareness and of low emotional awareness with latent factor scores for psychopathology, controlling for age, sex, and their interaction. All data and code are available at https://github.com/dgweissman/EMAware_P.

Results

Structure of Psychopathology. In initial models, PTSD symptoms did not load significantly onto the general factor in the bifactor model or to the internalizing factor in the correlated-factors model. Therefore, PTSD symptoms were excluded from final models. Consistent with prior work, the bifactor and correlated-factors model provided similar fit to the

data. Fit indices for the bi-factor model were: AIC = 2681.05, BIC = 2869.94, Sample adjusted BIC = 2708.20. Standardized factor loadings for the latent p-factor ranged from 0.47 to 0.93 (all p 's < .001). Fit indices for the correlated-factors model were: AIC = 2684.00, BIC = 2825.50, Sample adjusted BIC = 2676.76. Standardized factor loadings for the Internalizing (CDI, SCARED) latent factor were 0.99 and 0.67 respectively, both p 's < .001. Standardized factor loadings for the Externalizing (Aggressive Behaviors, Rule Breaking Behaviors, Attention Problem) latent factor ranged from 0.55 to 0.80, all p 's < .001. The internalizing and externalizing factors were correlated at .715, p < .001.

We extracted general factor scores from the bifactor model for further analyses because i) our primary interest was in psychopathology across the externalizing and internalizing spectrum, ii) both models fit the data similarly, and iii) we had some concerns about the reliability of the internalizing factor in the correlated factors model when estimated by only two indicators (Kline, 2015).

Age and Sex Differences in Emotional Awareness. Emotional awareness was marginally correlated with age ($r = .19, p = .056$), and males and females did not differ significantly in emotional awareness (*Cohen's d* = .10, $t = 0.49, p = .625$; Table 1). There was a significant age \times sex interaction in relation to emotional awareness ($B = 1.53, SE = 0.67, p = .024$). Low emotional awareness was positively associated with age in females ($r = .39, p = .004$) but not males ($r = -.05, p = .752$, Figure 1). Thus, emotional awareness decreased from childhood to adolescence for females but not males.

Age, Sex, and Psychopathology. The p-factor was marginally correlated with age ($r = .17, p = .059$). Males and females did not differ in the magnitude of the p-factor (*Cohen's d* = -.23, $t = -1.29, p = .200$). The age \times sex interaction in relation to the p-factor was marginal (B

$=.09, SE = .05, p = .068$).

Emotional awareness and psychopathology. Low emotional awareness was moderately and significantly correlated with the p-factor ($r = .49, p < .001$; Figure 1) in an unadjusted model. The association remained significant controlling for age, sex, and their interaction ($B = .04, SE = .008, p < .001$).

Discussion

Study 1 demonstrated that emotional awareness declines and psychopathology increases with age across middle childhood and adolescence among females, but not males, and that low emotional awareness was strongly associated with the p-factor. Although it may be intuitive to think that emotional awareness would improve across childhood and adolescence, we found the opposite. This finding replicates and extends prior work (e.g., van der Crujisen et al., 2019) by showing that worsening emotional awareness across this age window explains broad transdiagnostic psychopathology in females but not males. Additionally, the strong concurrent association between low emotional awareness and the p-factor suggests that it may be a transdiagnostic risk factor contributing to psychopathology. However, the cross-sectional nature of this study's design leaves the direction of this relationship unclear: low emotional awareness could also be a shared consequence of many forms of psychopathology symptoms. Similarly, even though these results suggest that increasing difficulties in emotional awareness in females might explain their increased symptoms of psychopathology during adolescence, the lack of a longitudinal design limits the application of mediation approaches (Maxwell & Cole, 2007).

Consequently, Study 2 aimed to replicate the findings of Study 1 in a larger dataset and extend them in two important ways. First, Study 2 utilized two waves of longitudinal data to examine whether low emotional awareness was not only concurrently associated with

psychopathology, but if it predicted change in psychopathology prospectively over time and whether it explained age-related differences in psychopathology among females. Second, Study 2 evaluated if childhood violence exposure was associated with low emotional awareness, and further if low emotional awareness was a mechanism linking violence exposure with psychopathology over time.

Study 2

Method

Participants. As reported on previously (Weissman et al., 2019), youth aged 8-16 years and a parent or guardian were recruited to participate in a study examining child violence exposure, emotion regulation, and psychopathology. A total of 262 youth aged 8-16 years were enrolled into the study. Exposure to maltreatment and other inclusion and exclusion criteria were assessed during the first study visit, along with several behavioral and self-report measures of emotion regulation. Youth and caregivers were recruited for participation at schools, after-school and prevention programs, adoption programs, food banks, shelters, parenting programs, medical clinics, and the general community in Seattle, WA between January 2015 and June 2017. Recruitment efforts were targeted at recruiting a sample with variation in exposure to violence. To do so, we recruited from neighborhoods with high levels of violent crime, from clinics that served a predominantly low-SES catchment area, and agencies that work with families who have been victims of violence (e.g., domestic violence shelters, programs for parents mandated to receive intervention by Child Protective Services). Inclusion criteria for the maltreated group included exposure to physical or sexual abuse or direct witnessing of domestic violence. Children in the control group were matched to children in the maltreated group on age, sex, and

handedness; inclusion criteria required an absence of exposure to maltreatment or other forms of significant interpersonal violence.

Exclusion criteria included $IQ < 80$, presence of pervasive developmental disorder, active psychotic symptoms or mania, active substance abuse, and presence of safety concerns. IQ was measured with the Wechsler Abbreviated Intelligence Scale (WASI) (Wechsler, 2011) in the first study visit. Pervasive developmental disorders were assessed via parent report. Of the 262 children enrolled in the first study visit, three were excluded from all analysis due to low IQ ($n=1$), presence of pervasive developmental disorder ($n=1$), and presence of psychotic symptoms and drug abuse ($n=1$). The total sample size for the present analysis was 259 children and adolescents. A longitudinal follow-up assessment was conducted approximately two years after baseline assessments ($M = 21.96$ months, $SD = 7.88$ months) to assess symptoms of psychopathology. A total of 198 children and adolescents (76.4%) completed the follow-up visit. Of the 65 children that dropped out, 44 were maltreated (28% attrition), and 21 were not (20% attrition). Attrition did not differ significantly between maltreated or control groups ($\chi^2 = 2.38, p = .123$).

All procedures were approved by the Institutional Review Board at the University of Washington. Written informed consent was obtained from legal guardians; children provided written assent. Maltreatment not previously reported to the relevant authorities was reported to Child Protective Services using standard clinical procedures. Children with active safety concerns were not enrolled in the study.

Measures. *Violence exposure.* For the purposes of this study, violence exposure included physical and sexual abuse, and chronic exposure to domestic violence. We used a multi-informant, multi-method approach for assessing exposure to violence. Children completed two

interviews with a trained member of our research team assessing child maltreatment experiences and exposure to interpersonal violence: the Childhood Experiences of Care and Abuse (CECA) Interview (Bifulco, Brown, & Harris, 1994) and the Violence Exposure Scale for Children-Revised (VEX-R) (Raviv et al., 2001; Raviv, Raviv, Shimoni, Fox, & Leavitt, 1999). The CECA assesses caregiving experiences, including physical and sexual abuse. We modified the interview to ask parallel questions about witnessing domestic violence (i.e., directly observing violence directed at a caregiver). Inter-rater reliability for maltreatment reports is excellent, and validation studies suggest high agreement between siblings on maltreatment reports (Bifulco, Brown, Lillie, & Jarvis, 1997). The VEX-R assesses the frequency of exposure to different forms of violence. Children are presented with a cartoon and caption depicting a child of the same sex witnessing a type of violence (e.g., “Chris sees a person slap another person really hard”) and experiencing that same type of violence (e.g., “A person slaps Chris really hard”). Children are then asked to report how frequently they have witnessed or experienced that type of violence (e.g., “How many times have you seen a person slap another person really hard?”; “How many times has a person slapped you really hard?”) on a Likert scale ranging from 0 (Never) to 3 (Lots of times). We added follow-up questions for each item that was endorsed to gather additional information (e.g., the perpetrator, age of onset). The VEX-R demonstrates good reliability and has been validated with children as young as second grade (Raviv et al., 2001, 1999).

Children also completed two self-report measures: the Childhood Trauma Questionnaire (CTQ) (Bernstein, Ahluvalia, Pogge, & Handelsman, 1997) and the UCLA PTSD Reaction Index (PTSD-RI) (Steinberg et al., 2004). The CTQ is a 28-item scale that assesses the frequency of maltreatment during childhood, including physical and sexual abuse. Validated thresholds for exposure to physical and sexual abuse (Walker et al., 1999) were applied here in evaluating

abuse exposure based on the CTQ. The CTQ has excellent psychometric properties including internal consistency, test-retest reliability, and convergent and discriminant validity with interviews and clinician reports of maltreatment (Bernstein et al., 1997, 1994). The PTSD-RI includes a trauma screen that assesses exposure to numerous traumatic events, including physical abuse, sexual abuse, and domestic violence and additionally assesses PTSD symptoms. The PTSD-RI has good internal consistency and convergent validity (Steinberg et al., 2013).

Caregivers completed three self-report measures: the Conflict Tactics Scale-Parent Child Version (CTS) (Straus, Hamby, Finkelhor, Moore, & Runyan, 1998), the Juvenile Victimization Questionnaire (JVQ) lifetime caregiver report (Finkelhor, Hamby, Ormrod, & Turner, 2005), and the caregiver version of the PTSD-RI. The CTS includes 22 items assessing caregiver responses to child disobedience or misbehavior in the past year. Caregivers indicate how frequently they have used each strategy (e.g., shook him/her) on a Likert scale ranging from 0 (This has never happened) to 6 (more than 20 times in the past year) and can also indicate if they have used the strategy in the past but not in the last year. The CTS has adequate reliability and good discriminant and construct validity (Straus et al., 1998). The JVQ includes 34 items assessing exposure to crime, child maltreatment, peer and sibling victimization, sexual victimization, and witnessing and indirect victimization and has excellent psychometric properties, including test-retest reliability and construct validity (Finkelhor et al., 2005). Caregivers endorsed whether their child had experienced each event in his/her lifetime. Caregivers also completed the trauma screen included in the PTSD-RI, described above. A trained interviewer followed up with the caregiver if the endorsed any form of abuse or domestic violence to gather additional information about the experience.

Children were classified as experiencing physical or sexual abuse if abuse was endorsed by the child (on the CECA interview, PTSD-RI trauma screen, or above the validated CTQ threshold) or parent (on the CTS, JVQ, or PTSD-RI trauma screen). A total of 100 children (38.6%) experienced physical or sexual abuse. Inter-rater reliability was fair to good for child and caregiver reports (82.0% agreement; kappa=0.62). Exposure to domestic violence (on the VEX-R interview or PTSD-RI trauma screen) was determined based on child report only. A total of 99 children (38.2%) reported witnessing domestic violence. Exposure to violence was coded dichotomously, with a 1 indicating that participants had been exposed to physical abuse, sexual abuse, or domestic violence, and a 0 indicating no exposure to those types of violence.

Emotional awareness. Emotional awareness was measured using the 8-item (e.g. “I have feelings that I can’t figure out”) Poor Emotional Awareness subscale of the Emotional Expressiveness Scale for Children (EESC), a well validated measure in children and adolescents (Penza-Clyve & Zeman, 2002). The scale had excellent reliability in this sample ($\alpha = .91$). Items are rated on a 5-point Likert scale from “not at all true” to “extremely true.” Higher scores indicate lower emotional awareness.

Psychopathology. Symptoms of psychopathology were measured using the same instruments as in Study 1, and the p-factor was defined using the same methods and criteria as in Study 1 (and as in Weissman et al., 2019). The CDI-2 was used to measure depressive symptoms. Anxiety symptoms were assessed with the SCARED. Attention problems, rule breaking behaviors, and aggressive behavior were assessed on the YSR and CBCL. Post-traumatic stress disorder symptoms were assessed with the PTSD-RI. We performed CFA to test a correlated-factors model and a bi-factor model. We binned scores on each indicator into deciles prior to CFA analyses. All CFA analyses were performed in MPlus version 8.1 using MLR. As

assessed by relative fit indices and factor loadings, both models fit the data well at the baseline assessment, with a relatively better fit for the bi-factor model. Latent factor scores for the p-factor from this model at baseline and follow-up were extracted for further analysis.

Analysis. First, linear regression was used to evaluate associations between age, sex, and their interaction with emotional awareness and between violence exposure and emotional awareness. We controlled for racial/ethnic minority status and income-to-needs ratio in these analyses, as these characteristics varied between violence-exposed and unexposed youth. Racial/ethnic minority status was a dichotomous variable indicating whether participants identified with a race or ethnicity other than white. Income-to-needs ratio was calculated by dividing the parent-reported family income by the poverty threshold for a family of that size as indicated by the U. S. Census Bureau (U.S. Census Bureau, 2019). Second, we investigated the association between age, sex, and their interaction with p-factor scores at baseline, and between violence exposure and p-factor scores at baseline, controlling for racial/ethnic minority status, and income-to-needs ratio.

Third, we evaluated the associations between emotional awareness and p-factor scores at baseline and at the follow-up, controlling for p-factor scores at baseline and the time elapsed from baseline to follow-up. Age, sex, their interaction, violence exposure, income-to-needs ratio, and racial/ethnic minority status were included as covariates in both analyses, due to their hypothesized association with emotional awareness and/or the p-factor. Controlling for the time elapsed between visits ensures that any differences in longitudinal change in the p-factor is not attributable to differences in the length of time since the previous measurement of the p-factor.

Fourth, a parallel mediation model with bootstrapped confidence intervals (10,000 iterations) was estimated using the lavaan package in R (Rosseel, 2012) to evaluate whether

emotional awareness mediated the association between violence exposure and the p-factor at follow-up and between the age and the p-factor at follow-up. A moderated mediation model was additionally evaluated to determine whether low emotional awareness mediated age-related increases in p-factor among females only. Because results of regression analyses revealed that the p-factor at baseline was the only covariate associated with the p-factor at follow-up, no other covariates were included in the mediation model. All data and code are available for download at https://github.com/dgweissman/EMAware_P.

Results

Structure of psychopathology. As described previously (Weissman et al., 2019), the correlated factor model and the bifactor model fit the data similarly, with a slightly better model fit for the bifactor model. In contrast to the models in Study 1, PTSD symptoms loaded significantly onto the p-factor in the bifactor model and onto the internalizing factor in the correlated factors model and were therefore included in the models. Factor scores for the p-factor were extracted for further analyses.

Age, sex, and emotional awareness. Age and low emotional awareness were positively correlated at baseline ($r = .18, p = .003$). Low emotional awareness did not differ significantly between males and females (*Cohen's d* = $-.10, t = -0.80, p = .425$). There was a significant age \times sex interaction in relation to emotional awareness ($B = .899, SE = 0.356, p = .015$), such that low emotional awareness was positively associated with age in females ($r = .35, p < .001$) but not males ($r = .03, p = .690$, Figure 1).

Childhood violence exposure and emotional awareness. Violence-exposed youth had higher scores on low emotional awareness than children and adolescents with no history of

violence exposure (*Cohen's d* = .57, *t* = 4.74, *p* < .001). This association remained robust and significant even after controlling for income-to-needs ratio, and racial minority status (*B* = 4.29, *SE* = 1.16, *p* < .001).

Age, sex, childhood violence exposure and psychopathology. There was a small but significant positive correlation between age and the p-factor at baseline (*r* = .15, *p* = .015). P-factor scores at baseline did not differ significantly between male and female adolescents (*Cohen's d* = .02, *t* = 0.19, *p* = .851), and the age × sex interaction in relation to p-factor at baseline was not significant (*B* = .0426, *SE* = .0430, *p* = .324).

Emotional awareness and psychopathology. Low emotional awareness was significantly correlated with the p-factor with a moderate effect size (*r* = .53, *p* < .001; Figure 1). This association remained robust and significant even after controlling for age, sex, their interaction, violence exposure, income-to-needs ratio, and racial minority status (*B* = .05, *SE* = .006, *p* < .001).

Low emotional awareness as a transdiagnostic mechanism. Low emotional awareness was significantly associated with the p-factor at follow-up, controlling for the p-factor at baseline, violence exposure, time between visits, age, sex, their interaction, racial minority status, and income-to-needs ratio (*B* = .02, *SE* = .007, *p* = .018; Table 2). When comparing the mediation and moderated mediation models, including the interaction between sex and age significantly improved model fit ($\chi^2 = 122.42$, *p* < .001), so the moderated mediation model was used. In both models, the indirect effect of violence exposure on the p-factor at follow-up via low emotional awareness did not include 0 (Unstandardized Estimate = .07, *SE* = .04, 95% *CI* = .008-.15), providing evidence of mediation of the effect of violence exposure on the p-factor by emotional awareness in all participants (Figure 2). The conditional indirect effect of age on the p-

factor at follow-up, with the path from age to emotional awareness conditional on sex, did not include 0, providing evidence of moderated mediation (Unstandardized Estimate = .02, $SE = .01$, 95% $CI = .001-.04$), such that lower emotional awareness only explained age-related increases in the p-factor in females (Figure 2).

Discussion

In a large longitudinal sample of children and adolescents, many of whom had been exposed to violence in the form of physical or sexual abuse or domestic violence, age was associated with low emotional awareness in females but not males, and low emotional awareness was positively associated with the p-factor at both baseline and follow-up. These results replicated the associations found in Study 1. When turning to other risk factors, we also found that exposure to violence was associated with lower emotional awareness. Finally, low emotional awareness predicted increases in the p-factor over time and mediated associations of both violence exposure and age with the p-factor, although the latter was true only for females. These results suggest that low emotional awareness may be a transdiagnostic mechanism linking childhood violence exposure with psychopathology and a mechanism of age-related increases in psychopathology among females.

General Discussion

We demonstrated in two studies of children and adolescents that emotional awareness declines from middle childhood through adolescence, is associated with transdiagnostic psychopathology concurrently, and predicts increased psychopathology over time. This suggests that low emotional awareness may be a factor that increases risk for the emergence of multiple forms of psychopathology across the internalizing and externalizing spectrum in adolescence. These longitudinal findings extend prior work associating low emotional awareness and specific

forms of psychopathology (Hendryx et al., 1991; Honkalampi et al., 2002; Taylor et al., 1990; van der Crujisen et al., 2019). Low emotional awareness also emerged as a factor explaining increases in transdiagnostic psychopathology as a function of adolescent development among females and exposure to violence. These findings suggesting that low emotional awareness may be a mechanism contributing to increased risk for emerging psychopathology during adolescence.

Low emotional awareness may contribute to psychopathology across the internalizing and externalizing spectrum in a number of ways. Difficulty assigning emotional labels to negative affect has been found to disrupt effective emotion regulation in prior studies (Barrett et al., 2001; Kalokerinos et al., 2019), and difficulties with emotion regulation are a well-established transdiagnostic risk factor for psychopathology (Aldao et al., 2010; Burns et al., 2010; McLaughlin, Aldao, et al., 2014a; McLaughlin, Aldao, Wisco, & Hilt, 2014b; McLaughlin & Nolen-Hoeksema, 2011b; Messman-Moore & Bhuptani, 2017). Thus, low emotional awareness may connect these two lines of work: Difficulties understanding what one is feeling may disrupt effective emotion regulation, leading to the emergence of multiple forms of psychopathology. However, low emotional awareness could also lead to psychopathology via a more social or interpersonal route. Low emotional awareness is associated with difficulties recognizing emotions in others (Collin et al., 2013; Grynberg et al., 2012; Nook, Lindquist, & Zaki, 2015). Difficulties identifying what other people are feeling may produce downstream issues in social support and interpersonal emotion regulation, which are associated with psychopathology (Hofmann, 2014; Marroquín, 2011; Williams, Morelli, Ong, & Zaki, 2018). Finally, assigning a label to emotional pictures results in reduced self-reported distress, psychophysiological responses, and amygdala reactivity in children, adolescents, and adults

(Burklund, David Creswell, Irwin, & Lieberman, 2014; Gee et al., 2015; Matthew D. Lieberman, Inagaki, Tabibnia, & Crockett, 2011; M.D. Lieberman et al., 2007; Tabibnia, Lieberman, & Craske, 2008). Given that heightened amygdala responses to affectively salient cues, particularly threatening stimuli, is associated with many forms of psychopathology in children and adolescents (Beesdo et al., 2009; Dotterer, Hyde, Swartz, Hariri, & Williamson, 2017; Monk et al., 2008; Swartz, Knodt, Radtke, & Hariri, 2015), this could be another pathway through which lower emotional awareness contributes to psychopathology risk. Thus, children and adolescents with low emotional awareness may lack a foundation for effective intrapersonal and interpersonal emotion regulation, putting them increased risk for psychopathology. However, specifically identifying the mechanistic processes through which these symptoms emerge remains an important area of future study.

In both studies, age was significantly related to low emotional awareness, but only in females. This finding suggests that the transition into adolescence may generate difficulties for females in identifying their emotions. These difficulties with emotional awareness, in turn, may contribute to increased risk for psychopathology. This is consistent with prior work suggesting that emotional awareness worsened across childhood and adolescence, and that low emotional awareness explained variance in the emergence of depression across age only in females (van der Crujisen et al., 2019). Building upon the ideas presented above, one potential explanation of these sex differences is that low emotional awareness may interfere with social support specifically in adolescent females. Indeed, longitudinal research in adolescents has found that emotional awareness predicts number of friends in females but not males (Rowell et al., 2014). Thus, entering adolescence without strong emotion identification and skills may lead females

(but not males) to struggle to build friendships, leading to difficulties finding social support, lower self-esteem and mood, and potentially the emergence of psychopathology.

We also found that childhood violence exposure was associated with low emotional awareness, and low emotional awareness mediated the association between childhood violence exposure and increases in the p-factor over time. Children exposed to violence and other forms of adversity have difficulties with labeling and differentiating between emotions (Pears & Fisher, 2005; Shenk et al., 2013), potentially due to stress-related disruptions in detecting internal bodily states (Barrett & Simmons, 2015; McEwen, 1998), chronic dysregulation in physiological stress response systems that may create a de-coupling between subjective affect and physiological activation (Carpenter et al., 2011; Gordis et al., 2010; Gunnar & Vazquez, 2001; Harkness et al., 2011; Heleniak et al., 2016; MacMillan et al., 2009; McLaughlin, Sheridan, et al., 2014; Trickett et al., 2010), a lack of consistent scaffolding and use of emotion language by parents (Denham et al., 1997; Pollak et al., 2000), or specialization in the detection of anger at the expense of differentiating between other types of negative emotions (Shenk et al., 2013). These findings extend prior work demonstrating that more normative forms of stressful life events, such as peer victimization, are associated with reductions in emotional awareness over time during adolescence (McLaughlin & Hatzenbuehler, 2009; McLaughlin et al., 2009). Thus, it may be that not only exposure to violence but a wider range of stressful experiences contribute to lower facility at categorizing emotions, leading to low emotional awareness, and worsening psychopathology over time. Further research is needed to examine this empirically.

Both studies identified similar hierarchical structures of psychopathology. However, in Study 1, PTSD symptoms did not load significantly on to the p-factor or the internalizing factor in the correlated factors model. This may reflect the unique characteristics of PTSD relative to

the other diagnoses included in the analyses, in that the existence of PTSD symptoms is contingent on the experiencing of a traumatic event. This speculation is supported by the finding in Study 1 that PTSD symptom severity was significantly correlated with low emotional awareness among youth who experienced trauma. The bifactor model and the correlated factor model both fit the data similarly in both Study 1 and Study 2. The precise structure of psychopathology in childhood and adolescence is a topic of rich, ongoing investigation in much larger datasets, which are better suited to answering that question (Castellanos-Ryan et al., 2016; Gomez et al., 2018; Laceulle et al., 2015; Martel et al., 2017; Murray et al., 2016; Neumann et al., 2016; Patalay et al., 2015; Schaefer et al., 2018; Snyder et al., 2017). However, both models converged in demonstrating that symptoms of psychopathology are highly comorbid in samples of children and adolescents and that low emotional awareness is transdiagnostically associated with higher psychopathology.

These findings point to the possibility that interventions aimed at improving emotional awareness could be effective at preventing the emergence of psychopathology, particularly in adolescent females and children who have experienced violence. Improving emotional awareness may also be useful as a treatment target for multiple forms of psychopathology. Better emotional awareness is thought to be an important mechanism through which existing effective interventions, such as the Unified Protocol, dialectical behavioral therapy, and mindfulness based therapies, operate (Ellard, Fairholme, Boisseau, Farchione, & Barlow, 2010; Hölzel et al., 2011; Lynch, Chapman, Rosenthal, Kuo, & Linehan, 2006). A better understanding of emotional awareness and its relation with violence exposure and psychopathology may contribute to improvements in existing evidence-based interventions.

This investigation had several strengths, including replication of findings across samples

(one normative sample and one that included children with high levels of exposure to violence and psychopathology), integration of multiple measures of violence exposure from both youth and parents, and longitudinal measures of psychopathology over time. However, two primary limitations constrain interpretation and suggest directions for future research. First, while the use of multiple reporters for quantifying violence exposure in Study 2 and psychopathology symptoms in both studies allays some concerns about associations being attributable to shared method variance, work utilizing behavioral and ecological (e.g., experience sampling) measures of emotional awareness would provide necessary additional support to our findings. Second, in Study 2, although violence-exposed and unexposed youth were matched on both age and sex, youth exposed to violence had significantly lower income and a higher proportion of racial/ethnic minority youth. These factors frequently co-occur, and all analyses controlled for income-to-needs ratio and racial minority status. However, potential confounds from this collinearity cannot be entirely ruled out. Future work in samples recruited specifically to examine interactions among violence exposure, race/ethnicity and socioeconomic status could help distinguish the unique and shared influence of these variables on emotional awareness and psychopathology.

Low emotional awareness is transdiagnostic mechanism underlying psychopathology in middle childhood and adolescence, as demonstrated by replication in two independent samples. Moreover, low emotional awareness appears to be a mechanism linking childhood violence exposure with psychopathology over time as well as age-related increases in psychopathology in females. Improving emotional awareness may therefore be an important target for early interventions designed to prevent psychopathology, particularly among adolescent females and youth exposed to violence.

Acknowledgements

This research was funded by the National Institute of Mental Health (R01-MH103291 to McLaughlin; K01-MH116325 to Miller), an Early Career Research Fellowship from the Jacobs Foundation (McLaughlin), and a OneMind Institute Rising Star Award (McLaughlin), and a National Science Foundation Graduate Research Fellowship (DGE1144152 to Nook).

References

- Achenbach, T. M. (1991). Manual for the child behavior checklist/4-18 and 1991 profile. In *Burlington VT*.
- Aldao, A., Nolen-Hoeksema, S., & Schweizer, S. (2010). Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review*.
<https://doi.org/10.1016/j.cpr.2009.11.004>
- Barrett, L. F. (2006). Solving the Emotion Paradox: Categorization and the Experience of Emotion. *Personality and Social Psychology Review*, *10*(1), 20–46.
https://doi.org/10.1207/s15327957pspr1001_2
- Barrett, L. F. (2017). The theory of constructed emotion: An active inference account of interoception and categorization. *Social Cognitive and Affective Neuroscience*.
<https://doi.org/10.1093/scan/nsw154>
- Barrett, L. F., Gross, J., Christensen, T. C., & Benvenuto, M. (2001). Knowing what you're feeling and knowing what to do about it: Mapping the relation between emotion differentiation and emotion regulation. *Cognition & Emotion*, *15*(6), 713–724.
<https://doi.org/10.1080/02699930143000239>
- Barrett, L. F., & Simmons, W. K. (2015). Interoceptive predictions in the brain. *Nature Reviews Neuroscience*. <https://doi.org/10.1038/nrn3950>
- Beesdo, K., Lau, J. Y. F., Guyer, A. E., McClure-Tone, E. B., Monk, C. S., Nelson, E. E., ... Pine, D. S. (2009). Common and distinct amygdala-function perturbations in depressed vs anxious adolescents. *Archives of General Psychiatry*, *66*(3), 275–285.
<https://doi.org/10.1001/archgenpsychiatry.2008.545>

- Bernstein, D. P., Ahluvalia, T., Pogge, D., & Handelsman, L. (1997). Validity of the childhood trauma questionnaire in an adolescent psychiatric population. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(3), 340–348.
<https://doi.org/10.1097/00004583-199703000-00012>
- Bernstein, D. P., Fink, L., Handelsman, L., Foote, J., Lovejoy, M., Wenzel, K., ... Ruggiero, J. (1994). Initial reliability and validity of a new retrospective measure of child abuse and neglect. *American Journal of Psychiatry*. <https://doi.org/10.1176/ajp.151.8.1132>
- Bifulco, A., Brown, G. W., & Harris, T. O. (1994). Childhood Experience of Care and Abuse (CECA): A Retrospective Interview Measure. *Journal of Child Psychology and Psychiatry*, 35(8), 1419–1435. <https://doi.org/10.1111/j.1469-7610.1994.tb01284.x>
- Bifulco, A., Brown, G. W., Lillie, A., & Jarvis, J. (1997). Memories of childhood neglect and abuse: Corroboration in a series of sisters. *Journal of Child Psychology and Psychiatry and Allied Disciplines*. <https://doi.org/10.1111/j.1469-7610.1997.tb01520.x>
- Birmaher, B., Brent, D. A., Chiappetta, L., Bridge, J., Monga, S., & Baugher, M. (1999). Psychometric properties of the screen for child anxiety related emotional disorders (SCARED): A replication study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38(10), 1230–1236. <https://doi.org/10.1097/00004583-199910000-00011>
- Birmaher, B., Khetarpal, S., Brent, D., Cully, M., Balach, L., Kaufman, J., & Neer, S. M. K. (1997). The Screen for Child Anxiety Related Emotional Disorders (SCARED): Scale construction and psychometric characteristics. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(4), 545–553. <https://doi.org/10.1097/00004583-199704000-00018>

- Burklund, L. J., David Creswell, J., Irwin, M. R., & Lieberman, M. D. (2014). The common and distinct neural bases of affect labeling and reappraisal in healthy adults. *Frontiers in Psychology*. <https://doi.org/10.3389/fpsyg.2014.00221>
- Burns, E. E., Jackson, J. L., & Harding, H. G. (2010). Child maltreatment, emotion regulation, and posttraumatic stress: The impact of emotional abuse. *Journal of Aggression, Maltreatment and Trauma*, *19*(8), 801–819.
<https://doi.org/10.1080/10926771.2010.522947>
- Carpenter, L. L., Shattuck, T. T., Tyrka, A. R., Geraciotti, T. D., & Price, L. H. (2011). Effect of childhood physical abuse on cortisol stress response. *Psychopharmacology*, *214*(1), 367–375. <https://doi.org/10.1007/s00213-010-2007-4>
- Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., ... Moffitt, T. E. (2014). The p factor: One general psychopathology factor in the structure of psychiatric disorders? *Clinical Psychological Science*.
<https://doi.org/10.1177/2167702613497473>
- Castellanos-Ryan, N., Briere, F. N., O'Leary-Barrett, M., Banaschewski, T., Bokde, A., Bromberg, U., ... The IMAGEN Consortium. (2016). The structure of psychopathology in adolescence and its common personality and cognitive correlates. *Journal of Abnormal Psychology*. <https://doi.org/10.1037/abn0000193>
- Collin, L., Bindra, J., Raju, M., Gillberg, C., & Minnis, H. (2013). Facial emotion recognition in child psychiatry: A systematic review. *Research in Developmental Disabilities*, *34*(5), 1505–1520. <https://doi.org/10.1016/j.ridd.2013.01.008>

- Craighead, W. E., Smucker, M. R., Craighead, L. W., & Ilardi, S. S. (1998). Factor analysis of the Children's Depression Inventory in a community sample. *Psychological Assessment, 10*(2), 156–165. <https://doi.org/10.1037/1040-3590.10.2.156>
- Deborde, A.-S., Vanwalleghem Maury, S., & Aitel, S. (2015). Emotion regulation in adolescents with conduct disorder and controls. *Encephale*.
<https://doi.org/10.1016/j.encep.2014.01.002>
- Denham, S. A., Mitchell-Copeland, J., Strandberg, K., Auerbach, S., & Blair, K. (1997). Parental contributions to preschoolers' emotional competence: Direct and indirect effects. *Motivation and Emotion*. <https://doi.org/10.1023/A:1024426431247>
- Dotterer, H. L., Hyde, L. W., Swartz, J. R., Hariri, A. R., & Williamson, D. E. (2017). Amygdala reactivity predicts adolescent antisocial behavior but not callous-unemotional traits. *Developmental Cognitive Neuroscience, 24*, 84–92.
<https://doi.org/10.1016/j.dcn.2017.02.008>
- Ellard, K. K., Fairholme, C. P., Boisseau, C. L., Farchione, T. J., & Barlow, D. H. (2010). Unified Protocol for the Transdiagnostic Treatment of Emotional Disorders: Protocol Development and Initial Outcome Data. *Cognitive and Behavioral Practice, 17*(1), 88–101. <https://doi.org/10.1016/j.cbpra.2009.06.002>
- Finkelhor, D., Hamby, S. L., Ormrod, R., & Turner, H. (2005). The Juvenile Victimization Questionnaire: Reliability, validity, and national norms. *Child Abuse and Neglect*.
<https://doi.org/10.1016/j.chiabu.2004.11.001>
- Frewen, P. A., Dozois, D. J. A., Neufeld, R. W. J., & Lanius, R. A. (2008). Meta-analysis of alexithymia in posttraumatic stress disorder. *Journal of Traumatic Stress*.
<https://doi.org/10.1002/jts.20320>

- Gatta, M., Facca, I., Colombo, E., Svanellini, L., Montagnese, S., & Schiff, S. (2014). Alexithymia, psychopathology and alcohol misuse in adolescence: A population based study on 3556 teenagers. *Neuroscience and Medicine*, *05*(01), 60–71.
<https://doi.org/10.4236/nm.2014.51009>
- Gee, D. G., McEwen, S. C., Forsyth, J. K., Haut, K. M., Bearden, C. E., Addington, J., ... Cannon, T. D. (2015). Reliability of an fMRI paradigm for emotional processing in a multisite longitudinal study. *Human Brain Mapping*. <https://doi.org/10.1002/hbm.22791>
- Gomez, R., Stavropoulos, V., Vance, A., & Griffiths, M. D. (2018). Re-evaluation of the Latent Structure of Common Childhood Disorders: Is There a General Psychopathology Factor (P-Factor)? *International Journal of Mental Health and Addiction*, 1–21.
<https://doi.org/10.1007/s11469-018-0017-3>
- Gordis, E. B., Feres, N., Olezeski, C. L., Rabkin, A. N., & Trickett, P. K. (2010). Skin Conductance Reactivity and Respiratory Sinus Arrhythmia Among Maltreated and Comparison Youth: Relations with Aggressive Behavior. *Journal of Pediatric Psychology*, *35*(5), 547–558. <https://doi.org/10.1093/jpepsy/jsp113>
- Grabe, H. J., Ruhrmann, S., Ettelt, S., Müller, A., Buhtz, F., Hochrein, A., ... Freyberger, H. J. (2006). Alexithymia in obsessive-compulsive disorder—Results from a family study. *Psychotherapy and Psychosomatics*. <https://doi.org/10.1159/000093954>
- Green, J. G., McLaughlin, K. A., Berglund, P. A., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2010). Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication I: Associations with first onset of DSM-IV disorders. *Archives of General Psychiatry*, *67*(2), 113–123.
<https://doi.org/10.1001/archgenpsychiatry.2009.186>

- Grynberg, D., Chang, B., Corneille, O., Maurage, P., Vermeulen, N., Berthoz, S., & Luminet, O. (2012). Alexithymia and the Processing of Emotional Facial Expressions (EFEs): Systematic Review, Unanswered Questions and Further Perspectives. *PLoS ONE*, 7(8). <https://doi.org/10.1371/journal.pone.0042429>
- Gunnar, M. R., & Vazquez, D. M. (2001). Low cortisol and a flattening of expected daytime rhythm: Potential indices of risk in human development. *Development and Psychopathology*, 13(3), 515–538.
- Harkness, K. L., Stewart, J. G., & Wynne-Edwards, K. E. (2011). Cortisol reactivity to social stress in adolescents: Role of depression severity and child maltreatment. *Psychoneuroendocrinology*, 36(2), 173–181. <https://doi.org/10.1016/j.psyneuen.2010.07.006>
- Heaven, P. C. L., Ciarrochi, J., & Hurrell, K. (2010). The distinctiveness and utility of a brief measure of alexithymia for adolescents. *Personality and Individual Differences*, 49(3), 222–227. <https://doi.org/10.1016/j.paid.2010.03.039>
- Heleniak, C., McLaughlin, K. A., Ormel, J., & Riese, H. (2016). Cardiovascular reactivity as a mechanism linking child trauma to adolescent psychopathology. *Biological Psychology*, 120, 108–119. <https://doi.org/10.1016/j.biopsycho.2016.08.007>
- Hendryx, M. S., Haviland, M. G., & Shaw, D. G. (1991). Dimensions of Alexithymia and Their Relationships to Anxiety and Depression. *Journal of Personality Assessment*. https://doi.org/10.1207/s15327752jpa5602_4
- Hofmann, S. G. (2014). Interpersonal Emotion Regulation Model of Mood and Anxiety Disorders. *Cognitive Therapy and Research*, 38(5), 483–492. <https://doi.org/10.1007/s10608-014-9620-1>

- Hölzel, B. K., Lazar, S. W., Gard, T., Schuman-Olivier, Z., Vago, D. R., & Ott, U. (2011). How does mindfulness meditation work? Proposing mechanisms of action from a conceptual and neural perspective. *Perspectives on Psychological Science*.
<https://doi.org/10.1177/1745691611419671>
- Honkalampi, K., Hintikka, J., Laukkanen, E., & Viinamäki, J. L. H. (2002). Alexithymia and Depression: A Prospective Study of Patients With Major Depressive Disorder. *Psychosomatics*. <https://doi.org/10.1176/appi.psy.42.3.229>
- Kalokerinos, E. K., Erbas, Y., Ceulemans, E., & Kuppens, P. (2019). Differentiate to Regulate: Low Negative Emotion Differentiation Is Associated With Ineffective Use but Not Selection of Emotion-Regulation Strategies. *Psychological Science*, *30*(6), 863–879.
<https://doi.org/10.1177/0956797619838763>
- Kashdan, T. B., Barrett, L. F., & McKnight, P. E. (2015). Unpacking Emotion Differentiation: Transforming Unpleasant Experience by Perceiving Distinctions in Negativity. *Current Directions in Psychological Science*, *24*(1), 10–16.
<https://doi.org/10.1177/0963721414550708>
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, *62*(6), 593–602.
<https://doi.org/10.1001/archpsyc.62.6.593>
- Keyes, K. M., Eaton, N. R., Krueger, R. F., McLaughlin, K. A., Wall, M. M., Grant, B. F., & Hasin, D. S. (2012). Childhood maltreatment and the structure of common psychiatric disorders. *British Journal of Psychiatry*. <https://doi.org/10.1192/bjp.bp.111.093062>

- Kline, R. B. (2015). *Principles and Practice of Structural Equation Modeling, Fourth Edition*. Guilford Publications.
- Kovacs, M. (1992). Children's Depression Inventory CDI Manual. *New York: Multi-Health Systems*. <https://doi.org/10.1103/PhysRevLett.101.208303>
- Kovacs, M. (2011). *Children's depression inventory (CDI2)*. North Tonawanda NY: Multi-Health Systems Inc.
- Laceulle, O. M., Vollebergh, W. A. M., & Ormel, J. (2015). The structure of psychopathology in adolescence: Replication of a general psychopathology factor in the TRAILS study. *Clinical Psychological Science*. <https://doi.org/10.1177/2167702614560750>
- Lahey, B. B., Applegate, B., Hakes, J. K., Zald, D. H., Hariri, A. R., & Rathouz, P. J. (2012). Is There a general factor of prevalent psychopathology during adulthood? *Journal of Abnormal Psychology*. <https://doi.org/10.1037/a0028355>
- Lane, R. D., & Schwartz, G. E. (1987). Levels of emotional awareness: A cognitive-developmental theory and its application to psychopathology. *The American Journal of Psychiatry*, *144*(2), 133–143. <https://doi.org/10.1176/ajp.144.2.133>
- Li, C. H. (2016). Confirmatory factor analysis with ordinal data: Comparing robust maximum likelihood and diagonally weighted least squares. *Behavior Research Methods*. <https://doi.org/10.3758/s13428-015-0619-7>
- Lieberman, Matthew D., Inagaki, T. K., Tabibnia, G., & Crockett, M. J. (2011). Subjective responses to emotional stimuli during labeling, reappraisal, and distraction. *Emotion*, *11*(3), 468–480. <https://doi.org/10.1037/a0023503>

- Lieberman, M.D., Eisenberger, N. I., Crockett, M. J., Tom, S. M., Pfeifer, J. H., & Way, B. M. (2007). Putting feelings into words. *Psychological Science*.
<https://doi.org/10.1111/j.1467-9280.2007.01916.x>
- Loas, G., Braun, S., Delhaye, M., & Linkowski, P. (2017). The measurement of alexithymia in children and adolescents: Psychometric properties of the Alexithymia Questionnaire for Children and the twenty-item Toronto Alexithymia Scale in different non-clinical and clinical samples of children and adolescents. *PLoS ONE*, *12*(5).
<https://doi.org/10.1371/journal.pone.0177982>
- Lumley, M. A., Neely, L. C., & Burger, A. J. (2007). The Assessment of Alexithymia in Medical Settings: Implications for Understanding and Treating Health Problems. *Journal of Personality Assessment*, *89*(3), 230–246. <https://doi.org/10.1080/00223890701629698>
- Lynch, T. R., Chapman, A. L., Rosenthal, M. Z., Kuo, J. R., & Linehan, M. M. (2006). Mechanisms of change in dialectical behavior therapy: Theoretical and empirical observations. *Journal of Clinical Psychology*, *62*(4), 459–480.
<https://doi.org/10.1002/jclp.20243>
- MacMillan, H. L., Georgiades, K., Duku, E. K., Shea, A., Steiner, M., Niec, A., ... Schmidt, L. A. (2009). Cortisol Response to Stress in Female Youths Exposed to Childhood Maltreatment: Results of the Youth Mood Project. *Biological Psychiatry*, *66*(1).
<https://doi.org/10.1016/j.biopsych.2008.12.014>
- Marroquín, B. (2011). Interpersonal emotion regulation as a mechanism of social support in depression. *Clinical Psychology Review*, *31*(8), 1276–1290.
<https://doi.org/10.1016/j.cpr.2011.09.005>

- Martel, M. M., Pan, P. M., Hoffmann, M. S., Gadelha, A., do Rosário, M. C., Mari, J. J., ...
Salum, G. A. (2017). A general psychopathology factor (P Factor) in children: Structural model analysis and external validation through familial risk and child global executive function. *Journal of Abnormal Psychology*. <https://doi.org/10.1037/abn0000205>
- Maxwell, S. E., & Cole, D. A. (2007). Bias in cross-sectional analyses of longitudinal mediation. *Psychological Methods*, *12*(1), 23–44. <https://doi.org/10.1037/1082-989X.12.1.23>
- McEwen, B. S. (1998). Stress, adaptation, and disease. Allostasis and allostatic load. *Annals of the New York Academy of Sciences*.
- McLaughlin, K. A., Aldao, A., Wisco, B. E., & Hilt, L. M. (2014a). Rumination as a transdiagnostic factor underlying transitions between internalizing symptoms and aggressive behavior in early adolescents. *Journal of Abnormal Psychology*.
<https://doi.org/10.1037/a0035358>
- McLaughlin, K. A., Aldao, A., Wisco, B. E., & Hilt, L. M. (2014b). Rumination as a transdiagnostic factor underlying transitions between internalizing symptoms and aggressive behavior in early adolescents. *Journal of Abnormal Psychology*.
<https://doi.org/10.1037/a0035358>
- McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2012). Childhood adversities and first onset of psychiatric disorders in a national sample of US adolescents. *Archives of General Psychiatry*.
<https://doi.org/10.1001/archgenpsychiatry.2011.2277>
- McLaughlin, K. A., & Hatzenbuehler, M. L. (2009). Mechanisms Linking Stressful Life Events and Mental Health Problems in a Prospective, Community-Based Sample of Adolescents. *Journal of Adolescent Health*. <https://doi.org/10.1016/j.jadohealth.2008.06.019>

- McLaughlin, K. A., Hatzenbuehler, M. L., & Hilt, L. M. (2009). Emotion dysregulation as a mechanism linking peer victimization to internalizing symptoms in adolescents. *Journal of Consulting and Clinical Psychology, 77*(5), 894–904.
<https://doi.org/10.1037/a0015760>
- McLaughlin, K. A., & Nolen-Hoeksema, S. (2011a). Rumination as a transdiagnostic factor in depression and anxiety. *Behaviour Research and Therapy.*
<https://doi.org/10.1016/j.brat.2010.12.006>
- McLaughlin, K. A., & Nolen-Hoeksema, S. (2011b). Rumination as a transdiagnostic factor in depression and anxiety. *Behaviour Research and Therapy.*
<https://doi.org/10.1016/j.brat.2010.12.006>
- McLaughlin, K. A., Sheridan, M. A., Alves, S., & Mendes, W. B. (2014). Child Maltreatment and Autonomic Nervous System Reactivity: Identifying Dysregulated Stress Reactivity Patterns by Using the Biopsychosocial Model of Challenge and Threat. *Psychosomatic Medicine, 76*(7), 538–546. <https://doi.org/10.1097/PSY.0000000000000098>
- Messman-Moore, T. L., & Bhuptani, P. H. (2017). A Review of the Long-Term Impact of Child Maltreatment on Posttraumatic Stress Disorder and Its Comorbidities: An Emotion Dysregulation Perspective. *Clinical Psychology: Science and Practice.*
<https://doi.org/10.1111/cpsp.12193>
- Monk, C. S., Telzer, E. H., Mogg, K., Bradley, B. P., Mai, X., Louro, H. M. C., ... Pine, D. S. (2008). Amygdala and Ventrolateral Prefrontal Cortex Activation to Masked Angry Faces in Children and Adolescents With Generalized Anxiety Disorder. *Archives of General Psychiatry, 65*(5), 568–576. <https://doi.org/10.1001/archpsyc.65.5.568>

- Murphy, J., Brewer, R., Catmur, C., & Bird, G. (2017). Interoception and psychopathology: A developmental neuroscience perspective. *Developmental Cognitive Neuroscience, 23*, 45–56. <https://doi.org/10.1016/J.DCN.2016.12.006>
- Murray, A. L., Eisner, M., & Ribeaud, D. (2016). The Development of the General Factor of Psychopathology ‘p Factor’ Through Childhood and Adolescence. *Journal of Abnormal Child Psychology*. <https://doi.org/10.1007/s10802-016-0132-1>
- Muthén, L. K., & Muthén, B. O. (2012). *Mplus User’s Guide* (7th ed.). Los Angeles, CA: Muthén & Muthén.
- Neumann, A., Pappa, I., Lahey, B. B., Verhulst, F. C., Medina-Gomez, C., Jaddoe, V. W., ... Tiemeier, H. (2016). Single Nucleotide Polymorphism Heritability of a General Psychopathology Factor in Children. *Journal of the American Academy of Child and Adolescent Psychiatry, 55*(12), 1038–1045. <https://doi.org/10.1016/j.jaac.2016.09.498>
- Nook, E. C., Lindquist, K. A., & Zaki, J. (2015). A new look at emotion perception: Concepts speed and shape facial emotion recognition. *Emotion (Washington, D.C.), 15*(5), 569–578. <https://doi.org/10.1037/a0039166>
- Nook, E. C., Sasse, S. F., Lambert, H. K., McLaughlin, K. A., & Somerville, L. H. (2017). Increasing verbal knowledge mediates development of multidimensional emotion representations. *Nature Human Behaviour, 1*(12), 881–889. <https://doi.org/10.1038/s41562-017-0238-7>
- Nook, E. C., Sasse, S. F., Lambert, H. K., McLaughlin, K. A., & Somerville, L. H. (2018). The nonlinear development of emotion differentiation: Granular emotional experience Is low in adolescence. *Psychological Science, 29*(8), 1346–1357. <https://doi.org/10.1177/0956797618773357>

- Nook, E. C., Stavish, C. M., Sasse, S. F., Lambert, H. K., Mair, P., McLaughlin, K. A., & Somerville, L. H. (in press). Charting the development of emotion comprehension and abstraction using observer-rated and linguistic measures. *Emotion*.
- O'Mahen, H. A., Karl, A., Moberly, N., & Fedock, G. (2015). The association between childhood maltreatment and emotion regulation: Two different mechanisms contributing to depression? *Journal of Affective Disorders, 174*, 287–295.
<https://doi.org/10.1016/J.JAD.2014.11.028>
- Patalay, P., Fonagy, P., Deighton, J., Belsky, J., Vostanis, P., & Wolpert, M. (2015). A general psychopathology factor in early adolescence. *British Journal of Psychiatry*.
<https://doi.org/10.1192/bjp.bp.114.149591>
- Pears, K., & Fisher, P. A. (2005). Emotion understanding and theory of mind among maltreated children in foster care: Evidence of deficits. *Development and Psychopathology*.
<https://doi.org/10.1017/S0954579405050030>
- Penza-Clyve, S., & Zeman, J. (2002). Initial Validation of the Emotion Expression Scale for Children (EESC). *Journal of Clinical Child and Adolescent Psychology*.
https://doi.org/10.1207/S15374424JCCP3104_12
- Pollak, S. D., Cicchetti, D., Hornung, K., & Reed, A. (2000). Recognizing emotion in faces: Developmental effects of child abuse and neglect. *Developmental Psychology*.
<https://doi.org/10.1037/0012-1649.36.5.679>
- Pollak, Seth D., & Sinha, P. (2002). Effects of early experience on children's recognition of facial displays of emotion. *Developmental Psychology*. <https://doi.org/10.1037/0012-1649.38.5.784>

- Raviv, A., Erel, O., Fox, N. A., Leavitt, L. A., Raviv, A., Dar, I., ... Greenbaum, C. W. (2001). Individual measurement of exposure to everyday violence among elementary schoolchildren across various settings. *Journal of Community Psychology*.
[https://doi.org/10.1002/1520-6629\(200103\)29:2<117::AID-JCOP1009>3.0.CO;2-2](https://doi.org/10.1002/1520-6629(200103)29:2<117::AID-JCOP1009>3.0.CO;2-2)
- Raviv, A., Raviv, A., Shimoni, H., Fox, N. A., & Leavitt, L. A. (1999). Children's Self-Report of Exposure to Violence and Its Relation to Emotional Distress. *Journal of Applied Developmental Psychology*, 20(2), 337–353. [https://doi.org/10.1016/S0193-3973\(99\)00020-9](https://doi.org/10.1016/S0193-3973(99)00020-9)
- Rieffe, C., Oosterveld, P., & Terwogt, M. M. (2006). An alexithymia questionnaire for children: Factorial and concurrent validation results. *Personality and Individual Differences*, 40(1), 123–133. <https://doi.org/10.1016/j.paid.2005.05.013>
- Rosseel, Y. (2012). lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical Software*, 48(2), 1–36. <https://doi.org/10.18637/jss.v048.i02>
- Rowell, H. C., Ciarrochi, J., Heaven, P. C. L., & Deane, F. P. (2014). The role of emotion identification skill in the formation of male and female friendships: A longitudinal study. *Journal of Adolescence*, 37(2), 103–111.
<https://doi.org/10.1016/j.adolescence.2013.11.005>
- Rybakowski, J., Ziółkowski, M., Zasadzka, T., & Brzeziński, R. (1988). High prevalence of alexithymia in male patients with alcohol dependence. *Drug and Alcohol Dependence*.
[https://doi.org/10.1016/0376-8716\(88\)90058-0](https://doi.org/10.1016/0376-8716(88)90058-0)
- Säkkinen, P., Kaltiala-Heino, R., Ranta, K., Haataja, R., & Joukamaa, M. (2007). Psychometric Properties of the 20-Item Toronto Alexithymia Scale and Prevalence of Alexithymia in a

Finnish Adolescent Population. *Psychosomatics*, 48(2), 154–161.

<https://doi.org/10.1176/appi.psy.48.2.154>

Salmon, K., O’Kearney, R., Reese, E., & Fortune, C.-A. (2016). The Role of Language Skill in Child Psychopathology: Implications for Intervention in the Early Years. *Clinical Child and Family Psychology Review*, 19(4), 352–367. <https://doi.org/10.1007/s10567-016-0214-1>

Satpute, A. B., & Lindquist, K. A. (2019). The Default Mode Network’s Role in Discrete Emotion. *Trends in Cognitive Sciences*, 0(0). <https://doi.org/10.1016/j.tics.2019.07.003>

Schaefer, J. D., Moffitt, T. E., Arseneault, L., Danese, A., Fisher, H. L., Houts, R., ... Caspi, A. (2018). Adolescent Victimization and Early-Adult Psychopathology: Approaching Causal Inference Using a Longitudinal Twin Study to Rule Out Noncausal Explanations. *Clinical Psychological Science*, 6(3), 352–371.

<https://doi.org/10.1177/2167702617741381>

Shenk, C. E., Putnam, F. W., & Noll, J. G. (2013). Predicting the accuracy of facial affect recognition: The interaction of child maltreatment and intellectual functioning. *Journal of Experimental Child Psychology*. <https://doi.org/10.1016/j.jecp.2012.08.007>

Snyder, H. R., Young, J. F., & Hankin, B. L. (2017). Strong Homotypic Continuity in Common Psychopathology-, Internalizing-, and Externalizing-Specific Factors Over Time in Adolescents. *Clinical Psychological Science*. <https://doi.org/10.1177/2167702616651076>

Steinberg, A. M., Brymer, M. J., Decker, K. B., & Pynoos, R. S. The University of California at Los Angeles Post-traumatic Stress Disorder Reaction Index. , 6 *Current Psychiatry Reports* § (2004).

- Steinberg, A. M., Brymer, M. J., Kim, S., Briggs, E. C., Ippen, C. G., Ostrowski, S. A., ... Pynoos, R. S. (2013). Psychometric Properties of the UCLA PTSD Reaction Index: Part I. *Journal of Traumatic Stress*. <https://doi.org/10.1002/jts.21780>
- Straus, M. A., Hamby, S. L., Finkelhor, D., Moore, D. W., & Runyan, D. (1998). Identification of child maltreatment with the parent-child Conflict Tactics Scales: Development and psychometric data for a national sample of American parents. *Child Abuse and Neglect*. [https://doi.org/10.1016/S0145-2134\(97\)00174-9](https://doi.org/10.1016/S0145-2134(97)00174-9)
- Swartz, J. R., Knodt, A. R., Radtke, S. R., & Hariri, A. R. (2015). A Neural Biomarker of Psychological Vulnerability to Future Life Stress. *Neuron*, *85*(3), 505–511. <https://doi.org/10.1016/J.NEURON.2014.12.055>
- Tabibnia, G., Lieberman, M. D., & Craske, M. G. (2008). The lasting effect of words on feelings: Words may facilitate exposure effects to threatening images. *Emotion*, *8*(3), 307–317. <https://doi.org/10.1037/1528-3542.8.3.307>
- Taylor, G. J., Parker, J. D. A., & Bagby, R. M. (1990). A preliminary investigation of alexithymia in men with psychoactive substance dependence. *American Journal of Psychiatry*. <https://doi.org/10.1176/ajp.147.9.1228>
- Trickett, P. K., Noll, J. G., Susman, E. J., Shenk, C. E., & Putnam, F. W. (2010). Attenuation of cortisol across development for victims of sexual abuse. *Development and Psychopathology*, *22*(1), 165–175. <https://doi.org/10.1017/S0954579409990332>
- U.S. Census Bureau. (2019). Poverty Thresholds. Retrieved from <https://www.census.gov/data/tables/time-series/demo/income-poverty/historical-poverty-thresholds.html>

- van der Crujisen, R., Murphy, J., & Bird, G. (2019). Alexithymic traits can explain the association between puberty and symptoms of depression and anxiety in adolescent females. *PLOS ONE*, *14*(1), e0210519. <https://doi.org/10.1371/journal.pone.0210519>
- van 't Wout, M., Aleman, A., Bermond, B., & Kahn, R. S. (2007). No words for feelings: Alexithymia in schizophrenia patients and first-degree relatives. *Comprehensive Psychiatry*. <https://doi.org/10.1016/j.comppsy.2006.07.003>
- Walker, E. A., Unutzer, J., Rutter, C., Gelfand, A., Saunders, K., VonKorff, M., ... Katon, W. (1999). Costs of health care use by women HMO members with a history of childhood abuse and neglect. *Archives of General Psychiatry*. <https://doi.org/10.1001/archpsyc.56.7.609>
- Wechsler, D. (2011). Wechsler Abbreviated Scale of Intelligence—Second Edition (WASI-II). In *Wechsler Abbreviated Scale of Intelligence—Second Edition (WASI-II)*. <https://doi.org/citeulike-article-id:6135820>
- Weissman, D. G., Bitran, D., Miller, A. B., Schaefer, J. D., Sheridan, M. A., & McLaughlin, K. A. (2019). Difficulties with emotion regulation as a transdiagnostic mechanism linking child maltreatment with the emergence of psychopathology. *Development and Psychopathology*, 1–17. <https://doi.org/10.1017/S0954579419000348>
- Williams, W. C., Morelli, S. A., Ong, D. C., & Zaki, J. (2018). Interpersonal emotion regulation: Implications for affiliation, perceived support, relationships, and well-being. *Journal of Personality and Social Psychology*, *115*(2), 224–254. <https://doi.org/10.1037/pspi0000132>

Table 1: Descriptive Statistics and Intercorrelations for Study 1

			Cohen's d								
	%	<i>n</i>	1	2	3	4	5	6	7	8	
Female	52	62	.07	.10	.21	.02	.12	-.25	-.30	-.31	
			Correlations								
	<i>N</i>	<i>M</i>	<i>SD</i>	1	2	3	4	5	6	7	8
1. Age	120	13.05	10.78	-							
2. Emotional awareness	104	25.00	3.44	.19	-						
3. Anxiety	102	13.79	12.00	.16	.62*	-					
4. Depression	100	6.29	6.56	.19	.51*	.62*	-				
5. PTSD	110	11.43	13.76	-.07	.08	.15	.14	-			
6. Aggression	120	6.78	4.58	-.15	.32*	.23*	.48*	.13	-		
7. Attention Problems	120	6.27	3.31	-.06	.36*	.26*	.57*	.20*	.60*	-	
8. Rule Breaking	120	3.60	2.99	.31*	.32*	.24*	.48*	.07	.48*	.35*	-

Note: * $p < .05$, N = number of participants with valid data on that measure, High scores on Emotional awareness indicate lower emotional awareness.

Table 2: Regression table of the relation between low emotional awareness and “p” at follow-up in Study 2

<i>Predictors</i>	P (follow-up)			
	<i>B</i>	<i>SE</i>	<i>β</i>	<i>p</i>
P (baseline)	.605*	.080	.608	<.001
Low emotional awareness	.0174*	.0073	.154	.018
Violence exposure	-.113	.138	-.064	.415
Age (Baseline)	.0750*	.0185	.224	<.001
Female	.0136	.0961	.008	.888
Age x Female	-.0588	.0371	-.087	.114
Time between visits (years)	.0783	.0744	.059	.294
Racial/ethnic minority status	-.081	.106	-.046	.443
Income-to-needs ratio	-.0046	.0220	-.015	.833
Observations	175			
R ² / adjusted R ²	0.528 / 0.502			

* p < .05

Figure 1: Scatterplots of relations between sex, age, low emotional awareness, and the p-factor




Figure 1: Low emotional awareness was measured with different instruments in studies 1 and 2. For ease of interpretation and visualization purposes, both scales were centered on their minimum value. Higher scores indicate lower emotional awareness. M = male, F = female, Age is in years.

Figure 2: Parallel mediation/moderated mediation of violence exposure and age (among females) on the p-factor through low emotional awareness in Study 2




Figure 2: All coefficients are standardized. Solid lines indicate significant paths. Dotted lines indicate nonsignificant paths. Main effects sex (female = 1) on emotional awareness and p-factor at follow-up were not significant and are not displayed but were also included in the model.